

**This document was generated by the Reference Group to the United Nations on HIV and Injecting Drug Use.**

**The data included in this report reflect the results of a systematic search and review of data on a range of indicators reflecting the extent of services provided to people who inject drugs.**

**These data have been searched, reviewed and extracted according to decision rules developed by the Reference Group.**

**Some data have been adjusted to facilitate estimates of standard indicators across countries.**

**Details of the process for searching, grading and including data, along with their interpretation, are detailed in:**

Mathers B, Degenhardt L, Ali H, Wiessing L, Hickman M, Mattick R, et al. HIV prevention, treatment and care for people who inject drugs: A systematic review of global, regional and national coverage. *The Lancet* 2010. DOI:10.1016/S0140-6736(10)60232-2.

**For more information on the Reference Group, please go to: [www.IDURefGroup.Com](http://www.IDURefGroup.Com)**

Section	Page
Estimated number of people who inject drugs and number of injectors who are living with HIV	4
<b>1. Needle and syringe programs (NSPs)</b>	<b>5</b>
NSP 1: Availability of NSPs	
NSP 2: Number of NSP sites	
NSP 3: Number of pharmacy sites where needles and syringes are available for purchase	
NSP 4: Total number of sites where needles and syringes are available	
NSP 5: Number of individual IDUs in contact with NSPs in a 12 month period	
NSP 6: Number of NSP occasions of service (client contacts) in a 12 month period	
NSP 7: Number of needles and syringes distributed by NSPs in a 12 month period	
<b>2. Opioid substitution programs (OST)</b>	<b>10</b>
OST 1: Availability of opioid substitution treatment (OST)	
OST 2: Availability of methadone maintenance treatment (MMT)	
OST 3: Availability of buprenorphine maintenance treatment (BMT)	
OST 4: Availability of other forms of agonist OST	
OST 5: Dose and duration of MMT	
OST 6: Dose and duration of BMT	
OST 7: Total number of all sites providing OST	
OST 8: Number of sites providing MMT	
OST 9: Number of sites providing BMT	
OST 10: Number of sites providing other forms of agonist OST	
OST 11: Number of all clients in any OST at a single point in time	
OST 12: Number of all clients in MMT at a single point in time	
OST 13: Number of all clients in BMT at a single point in time	
OST 14: Number of all clients in other forms of OST at a single point in time	
OST 15: Number of IDUs in any OST at a single point in time	
OST 16: Number of IDUs in MMT at a single point in time	
OST 17: Number of IDUs in BMT at a single point in time	
OST 18: Number of IDUs in other forms of OST at a single point in time	
OST 19: Number of all clients receiving any OST in a 12 month period	
OST 20: Number of all clients receiving MMT in a 12 month period	
OST 21: Number of all clients receiving BMT in a 12 month period	
OST 22: Number of all clients receiving other forms of OST in a 12 month period	
OST 23: Number of IDUs receiving any OST in a 12 month period	
OST 24: Number of IDUs receiving MMT in a 12 month period	

Section	Page
OST 25: Number of IDUs receiving BMT in a 12 month period	
OST 26: Number of IDUs receiving other forms of OST in a 12 month period	
<b>3. Other Drug Treatment (ODT)</b>	<b>25</b>
<hr/>	
ODT 1: Detoxification	
ODT 1.1: Number of sites providing detoxification	
ODT 1.2: Number of IDUs receiving detoxification	
ODT 1.3: Number of all clients receiving detoxification in a 12 month period and/or at a single point in time	
ODT 2: Residential Rehabilitation	
ODT 2.1: Number of sites providing residential rehabilitation	
ODT 2.2: Number of IDUs in residential rehabilitation	
ODT 2.3: Number of all clients in residential rehabilitation	
ODT 3: Outpatient psychosocial support, counselling and psychotherapy	
ODT 3.1: Number of sites providing outpatient psychosocial support, counselling or psychotherapy	
ODT 3.2: Number of IDUs receiving outpatient psychosocial support, counselling or psychotherapy treatment	
ODT 3.3: Number of all clients receiving outpatient psychosocial support, counselling or psychotherapy treatment	
ODT 4: Peer-based support groups	
ODT 4.1: Number of sites providing peer-based support groups	
ODT 4.2: Number of IDUs in peer-based support groups	
ODT 4.3: Number of all clients in peer-based support groups	
ODT 5: Drug treatment (treatment type not specified)	
ODT 5.1: Number of sites providing unspecified drug treatment	
ODT 5.2: Number of IDUs receiving unspecified drug treatment	
ODT 5.3: Number of all clients receiving unspecified drug treatment	
ODT 6: Drug treatment in prisons	
ODT 6.1: Number of drug treatment sites located in prisons	
ODT 6.2: Number of IDUs receiving drug treatment in prisons	
ODT 6.3: Number of all clients receiving drug treatment in prisons	
ODT 7: Compulsory drug treatment	
ODT 7.1: Number of compulsory detention sites for drug use	
ODT 7.2: Number of IDUs in compulsory detention sites for drug use	
ODT 7.3: Number of all drug users in compulsory detention sites for drug use	
<b>4. HIV testing and counselling (T&amp;C)</b>	<b>40</b>
<hr/>	
T&C 1: Availability of HIV T&C	
T&C 2: Number of sites providing HIV T&C that specifically targets IDUs	

<b>Section</b>	<b>Page</b>
T&C 3: Number of T&C sites providing HIV T& to the general population and also accessible to IDUs	
T&C 4: Number of individual IDUs who were tested for HIV in a 12 month period	
T&C 5: Percentage of IDUs who were tested for HIV in a 12 month period and who know the result	
<b>5. Anti-retro-viral (ARV) treatment</b>	<b>43</b>
<hr/>	
ARV 1: Number of sites providing ARV treatment	
ARV 2: Number of IDUs receiving ARVs at a single point in time	
ARV 3: Restrictions on IDUs' access to ARVs	
<b>6. Condoms</b>	<b>46</b>
<hr/>	
Condoms 1: Availability of condom programs targeting IDUs	
Condoms 2: Number of sites distributing condoms to IDUs	
Condoms 3: Number of individual IDUs receiving condoms	
Condoms 4: Number of condoms distributed to IDUs	

## Azerbaijan (Eastern Europe)

### Estimated number of people who inject drugs and number of injectors who are living with HIV

<b>Estimated prevalence of IDU among general population aged 15-64 years (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	3.83	5.21	6.81
<b>Year of IDU prevalence estimate</b>	2006		
<b>Reference for IDU prevalence estimate</b>	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.  Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
<b>Estimated prevalence of HIV among people who inject (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	2	13	24
<b>Year of HIV prevalence among IDU estimate</b>	2004		
<b>Reference for HIV prevalence among IDU estimate</b>	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
<b>General population 15-64 years of age in 2008</b>	5 990 102 (source: UN Population Division, 2009)		
<b>Estimated IDU population in 2008</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	229 500	312 000	407 500
<b>Estimated number of IDUs living with HIV in 2008</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	4590	40 560	97 800

## 1. Needle and syringe programs (NSPs)

### NSP 1: Availability of NSPs

Are NSPs available?	Yes
Comments	
Most recent year of documentation	2009
Reference	See references for other NSP indicators
When were NSPs first introduced?	1999
Reference	Zabransky, T. (Research and Development Manager), Center for Addictology Charles University / 1st Medical Faculty / Psychiatric Clinic Czech Republic,. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Czech Republic, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 18 May 2009.
Are needles and syringes available for purchase from pharmacies?	
Most recent year of documentation	
Reference	

### NSP 2: Number of NSP sites

Data reported	12 sites 14 NSPs		
Year of reported data	2008 2007		
Reference	WHO, UNAIDS., et al. (2009). Towards universal access: Scaling up priority HIV/AIDS interventions in the health sector - progress report 2009. Geneva, WHO.  WHO and UNAIDS (2008). Progress on implementing the Dublin Declaration on partnership to fight HIV/AIDS in Europe & Central Asia WHO.		
Adjustments	Figures taken as range		
Final/Adjusted data (to be used as indicator numerator)	12 - 14		
IDU prevalence estimate (%)	Low	Mid	High
	3.83	5.21	6.81
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.  Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
General population 15-64 years of age at year of reported data	5 990 102 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	229 500	312 000	407 500
INDICATOR: Number of NSP sites per 1000 IDUs	Low	Mid	High
	0.03	0.04	0.06

## NSP 3: Number of pharmacy sites where needles and syringes are available for purchase

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Number of pharmacy sites per 1000 IDUs	Low	Mid	High

## NSP 4: Total number of sites where needles and syringes are available

Total number (NSP sites + Pharmacy sites)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Total number of sites per 1000 IDUs	Low	Mid	High

## NSP 5: Number of individual IDUs in contact with NSPs in a 12 month period

Data reported	NSP present but number not known		
Year of reported data	2007		
Reference	WHO and UNAIDS (2008). Progress on implementing the Dublin Declaration on partnership to fight HIV/AIDS in Europe & Central Asia WHO.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	3.83	5.21	6.81
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.  Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
General population 15-64 years of age at year of reported data	5 869 830 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	225 000	305 500	399 500
INDICATOR: Percentage of IDUs in contact with NSPs in a 12 month period	Low	Mid	High
	NK	NK	NK

## NSP 6: Number of NSP occasions of service (client contacts) in a 12 month period

Data reported	NSP present but number of occasions of service not known		
Year of reported data	2007		
Reference	WHO and UNAIDS (2008). Progress on implementing the Dublin Declaration on partnership to fight HIV/AIDS in Europe & Central Asia WHO.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	3.83	5.21	6.81
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.  Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
General population 15-64 years of age at year of reported data	5 869 830 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	225 000	305 500	399 500
INDICATOR: Number of NSP occasions of service (client contacts) in a 12 month period per IDU	Low	Mid	High
	NK	NK	NK

## NSP 7: Number of needles and syringes distributed by NSPs in a 12 month period

Data reported	NSP present but number not known		
Year of reported data	2007		
Reference	WHO and UNAIDS (2008). Progress on implementing the Dublin Declaration on partnership to fight HIV/AIDS in Europe & Central Asia WHO.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	3.83	5.21	6.81
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.  Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
General population 15-64 years of age at year of reported data	5 869 830 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	225 000	305 500	399 500
INDICATOR: Total number of needles/syringes distributed in a 12 month period per IDU	Low	Mid	High
	NK	NK	NK

**NSP: All documents reviewed**

Aceijas, C., M. Hickman, et al. (2007). "Access and coverage of needle and syringe programmes (NSP) in Central and Eastern Europe and Central Asia.[see comment]." *Addiction* 102(8): 1244-50.

Afet Naibova Azerbaijan (2009). Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Reference Group to the UN on HIV and Injecting Drug Use.

AVERT. (2008). "HIV and AIDS in Russian Eastern Europe & Central Asia." Retrieved 30 November, 2008, from <http://www.avert.org/aids-russia.htm>

International Harm Reduction Development Program (2008). *Harm reduction developments 2008*. New York.

Penitentiary Services, Main Medical Department, et al. (2008). *HIV/AIDS Knowledge Attitude and Practice (KAP): "Anonymous Survey among Prisoners and Prison Staff in the Penitentiary System of the Republic of Azerbaijan"*.

Tomas Zabransky Research and Development Manager Center for Addictology Charles University / 1st Medical Faculty / Psychiatric Clinic Czech Republic (2009). Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Reference Group to the UN on HIV and Injecting Drug Use.

WHO and UNAIDS (2008). *Progress on implementing the Dublin Declaration on partnership to fight HIV/AIDS in Europe & Central Asia* WHO.

WHO, UNAIDS., et al. (2009). *Towards universal access: Scaling up priority HIV/AIDS interventions in the health sector - progress report 2009*. Geneva, WHO.

Zabransky, T. (Research and Development Manager), Center for Addictology Charles University / 1st Medical Faculty / Psychiatric Clinic Czech Republic,. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Czech Republic, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 18 May 2009.

**2. Opioid substitution treatment (OST)****OST 1: Availability of opioid substitution treatment (OST)**

Are any forms of agonist OST available?	Yes
Comments	
Most recent year of documentation	2009
Reference	See references for other OST indicators
When was agonist OST first available?	
Reference	

**OST 2: Availability of methadone maintenance treatment (MMT)**

Is MMT available?	Yes
Comments	
Most recent year of documentation	2009
Reference	See references for other OST indicators
When was MMT first available?	
Reference	

**OST 3: Availability of buprenorphine maintenance treatment (BMT)**

Is BMT available?	No
Comments	
Most recent year of documentation	2009
Reference	Naibova, A. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Azerbaijan, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 16 May 2009.  Tomas Zabransky Research and Development Manager Center for Addictology Charles University / 1st Medical Faculty / Psychiatric Clinic Czech Republic (2009). Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Reference Group to the UN on HIV and Injecting Drug Use.
When was BMT first available?	
Reference	

**OST 4: Availability of other forms of agonist OST**

Are other forms of OST available?	
Comments	
Most recent year of documentation	
Reference	
When were other forms of OST first available?	
Reference	

## OST 5: Dose and duration of MMT

Average/standard maintenance dose of MMT	60 - 70 mg
Comments	
Year	2009
Reference	World Health Organisation (2009). Global health ATLAS data.
Average/standard duration of MMT (range)	
Comments	
Year	
Reference	

## OST 6: Dose and duration of BMT

Average/standard maintenance dose of BMT	
Comments	
Year	
Reference	
Average/standard duration of BMT (range)	
Comments	
Year	
Reference	

## OST 7: Total number of all sites providing OST

Data reported	2 MMT sites 0 BMT sites		
Year of reported data	2009		
Reference	WHO, UNAIDS., et al. (2009). Towards universal access: Scaling up priority HIV/AIDS interventions in the health sector - progress report 2009. Geneva, WHO. UNODC (2009). Status of OST provision and plans for 2010.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	2		
IDU prevalence estimate (%)	Low	Mid	High
	3.83	5.21	6.81
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745. Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
General population 15-64 years of age at year of reported data	6 104 384 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	234 000	318 000	415 500
INDICATOR: Number of OST sites per 1000 IDUs	Low	Mid	High
	0.00	0.01	0.01

## OST 8: Number of sites providing MMT

Data reported	2 MMT sites		
Year of reported data	2009		
Reference	UNODC (2009). Status of OST provision and plans for 2010.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	2		
IDU prevalence estimate (%)	Low	Mid	High
	3.83	5.21	6.81
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.  Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
General population 15-64 years of age at year of reported data	6 104 384 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	234 000	318 000	415 500
INDICATOR: Number of MMT sites per 1000 IDUs	Low	Mid	High
	0.00	0.01	0.01

## OST 9: Number of sites providing BMT

Data reported	BMT not available		
Year of reported data	2009		
Reference	Naibova, A. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Azerbaijan, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 16 May 2009.  Tomas Zabransky Research and Development Manager Center for Addictology Charles University / 1st Medical Faculty / Psychiatric Clinic Czech Republic (2009). Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Reference Group to the UN on HIV and Injecting Drug Use.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	3.83	5.21	6.81
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.  Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
General population 15-64 years of age at year of reported data	6 104 384 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	234 000	318 000	415 500
INDICATOR: Number of BMT sites per 1000 IDUs	Low	Mid	High
	0	0	0

## OST 10: Number of sites providing other forms of agonist OST

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Number of sites providing other forms of OST per 1000 IDUs	Low	Mid	High

## OST 11: Number of all clients in any OST at a single point in time

<b>Data reported</b>	100-110 clients in MMT 0 clients in BMT		
<b>Year of reported data</b>	2008 2009		
<b>Reference</b>	<p>UNODC (2009). Status of OST provision and plans for 2010.</p> <p>Tomas Zabransky Research and Development Manager Center for Addictology Charles University / 1st Medical Faculty / Psychiatric Clinic Czech Republic (2009). Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Reference Group to the UN on HIV and Injecting Drug Use.</p> <p>Naibova, A. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Azerbaijan, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 16 May 2009.</p> <p>World Health Organisation (2009). Global health ATLAS data.</p>		
<b>Adjustments</b>			
<b>Final/Adjusted data (to be used as indicator numerator)</b>	100-110		
<b>IDU prevalence estimate (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	3.83	5.21	6.81
<b>Reference</b>	<p>Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.</p> <p>Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.</p>		
<b>General population 15-64 years of age at year of reported data</b>	6 104 384 (source: UN Population Division, 2009)		
<b>Estimated number of IDUs at year of reported data (to be used as indicator denominator)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	234 000	318 000	415 500
<b>INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on any OST at census date : 100 IDUs</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	0.02	0.03	0.05

## OST 12: Number of all clients in MMT at a single point in time

<b>Data reported</b>	100 clients enrolled in MMT at 1 Dec 2008 110 clients enrolled in MMT in 2008 110 patients on MMT		
<b>Year of reported data</b>	2008 2009		
<b>Reference</b>	UNODC (2009). Status of OST provision and plans for 2010.  Tomas Zabransky Research and Development Manager Center for Addictology Charles University / 1st Medical Faculty / Psychiatric Clinic Czech Republic (2009). Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Reference Group to the UN on HIV and Injecting Drug Use.  Naibova, A. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Azerbaijan, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 16 May 2009.  World Health Organisation (2009). Global health ATLAS data.		
<b>Adjustments</b>	Figures taken as a range		
<b>Final/Adjusted data (to be used as indicator numerator)</b>	100-110		
<b>IDU prevalence estimate (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	3.83	5.21	6.81
<b>Reference</b>	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.  Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
<b>General population 15-64 years of age at year of reported data</b>	6 104 384 (source: UN Population Division, 2009)		
<b>Estimated number of IDUs at year of reported data (to be used as indicator denominator)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	234 000	318 000	415 500
<b>INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on MMT at census date : 100 IDUs</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	0.02	0.03	0.05

## OST 13: Number of all clients in BMT at a single point in time

Data reported	BMT not available		
Year of reported data	2009		
Reference	Naibova, A. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Azerbaijan, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 16 May 2009.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	3.83	5.21	6.81
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.  Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
General population 15-64 years of age at year of reported data	6 104 384 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	234 000	318 000	415 500
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on BMT at census date : 100 IDUs	Low	Mid	High
	0	0	0

## OST 14: Number of all clients in other forms of OST at a single point in time

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on other forms of OST at census date : 100 IDUs	Low	Mid	High

## OST 15: Number of IDUs in any OST at a single point in time

<b>Data reported</b>	OST available but number of IDU on OST not known		
<b>Year of reported data</b>	2008		
<b>Reference</b>	<p>UNODC (2009). Status of OST provision and plans for 2010.</p> <p>Tomas Zabransky Research and Development Manager Center for Addictology Charles University / 1st Medical Faculty / Psychiatric Clinic Czech Republic (2009). Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Reference Group to the UN on HIV and Injecting Drug Use.</p> <p>Naibova, A. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Azerbaijan, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 16 May 2009.</p>		
<b>Adjustments</b>			
<b>Final/Adjusted data (to be used as indicator numerator)</b>	NK		
<b>IDU prevalence estimate (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	3.83	5.21	6.81
<b>Reference</b>	<p>Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.</p> <p>Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.</p>		
<b>General population 15-64 years of age at year of reported data</b>	5 990 102 (source: UN Population Division, 2009)		
<b>Estimated number of IDUs at year of reported data (to be used as indicator denominator)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	229 500	312 000	407 500
<b>INDICATOR: Percentage of IDUs in OST at a single point in time</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	NK	NK	NK

## OST 16: Number of IDUs in MMT at a single point in time

Data reported	MMT available but number of IDUs not known		
Year of reported data	2008		
Reference	<p>UNODC (2009). Status of OST provision and plans for 2010.</p> <p>Tomas Zabransky Research and Development Manager Center for Addictology Charles University / 1st Medical Faculty / Psychiatric Clinic Czech Republic (2009). Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Reference Group to the UN on HIV and Injecting Drug Use.</p> <p>Naibova, A. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Azerbaijan, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 16 May 2009.</p>		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	3.83	5.21	6.81
Reference	<p>Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.</p> <p>Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.</p>		
General population 15-64 years of age at year of reported data	5 990 102 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	229 500	312 000	407 500
INDICATOR: Percentage of IDUs in MMT at a single point in time	Low	Mid	High
	NK	NK	NK

## OST 17: Number of IDUs in BMT at a single point in time

Data reported	BMT not available		
Year of reported data	2009		
Reference	<p>Naibova, A. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Azerbaijan, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 16 May 2009.</p>		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	3.83	5.21	6.81
Reference	<p>Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.</p> <p>Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.</p>		
General population 15-64 years of age at year of reported data	6 104 384 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	234 000	318 000	415 500
INDICATOR: Percentage of IDUs in BMT at a single point in time	Low	Mid	High
	0	0	0

## OST 18: Number of IDUs in other forms of OST at a single point in time

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Percentage of IDUs in other forms of OST at a single point in time	Low	Mid	High

## OST 19: Number of all clients receiving any OST in a 12 month period

Data reported	OST available but number of clients on OST not known		
Year of reported data	2008		
Reference	<p>UNODC (2009). Status of OST provision and plans for 2010.</p> <p>Tomas Zabransky Research and Development Manager Center for Addictology Charles University / 1st Medical Faculty / Psychiatric Clinic Czech Republic (2009). Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Reference Group to the UN on HIV and Injecting Drug Use.</p> <p>Naibova, A. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Azerbaijan, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 16 May 2009.</p>		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	3.83	5.21	6.81
Reference	<p>Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.</p> <p>Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.</p>		
General population 15-64 years of age at year of reported data	5 990 102 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	229 500	312 000	407 500
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on any OST in a 12 month period : 100 IDUs	Low	Mid	High
	NK	NK	NK

## OST 20: Number of all clients receiving MMT in a 12 month period

Data reported	MMT available but number of clients not known		
Year of reported data	2008		
Reference	<p>UNODC (2009). Status of OST provision and plans for 2010.</p> <p>Tomas Zabransky Research and Development Manager Center for Addictology Charles University / 1st Medical Faculty / Psychiatric Clinic Czech Republic (2009). Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Reference Group to the UN on HIV and Injecting Drug Use.</p> <p>Naibova, A. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Azerbaijan, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 16 May 2009.</p>		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	3.83	5.21	6.81
Reference	<p>Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.</p> <p>Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.</p>		
General population 15-64 years of age at year of reported data	5 990 102 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	229 500	312 000	407 500
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on MMT in a 12 month period : 100 IDUs	Low	Mid	High
	NK	NK	NK

## OST 21: Number of all clients receiving BMT in a 12 month period

Data reported	BMT not available		
Year of reported data	2009		
Reference	<p>Naibova, A. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Azerbaijan, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 16 May 2009.</p>		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	3.83	5.21	6.81
Reference	<p>Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.</p> <p>Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.</p>		
General population 15-64 years of age at year of reported data	6 104 384 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	234 000	318 000	415 500
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on BMT in a 12 month period : 100 IDUs	Low	Mid	High
	0	0	0

## OST 22: Number of all clients receiving other forms of OST in a 12 month period

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on other forms of OST in a 12 month period : 100 IDUs	Low	Mid	High

## OST 23: Number of IDUs receiving any OST in a 12 month period

Data reported	OST available but number of IDU on OST not known		
Year of reported data	2008		
Reference	<p>UNODC (2009). Status of OST provision and plans for 2010.</p> <p>Tomas Zabransky Research and Development Manager Center for Addictology Charles University / 1st Medical Faculty / Psychiatric Clinic Czech Republic (2009). Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Reference Group to the UN on HIV and Injecting Drug Use.</p> <p>Naibova, A. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Azerbaijan, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 16 May 2009.</p>		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	3.83	5.21	6.81
Reference	<p>Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.</p> <p>Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.</p>		
General population 15-64 years of age at year of reported data	5 990 102 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	229 500	312 000	407 500
INDICATOR: Percentage of IDUs receiving OST in a 12 month period	Low	Mid	High
	NK	NK	NK

## OST 24: Number of IDUs receiving MMT in a 12 month period

Data reported	MMT available but number of IDUs not known		
Year of reported data	2008		
Reference	<p>UNODC (2009). Status of OST provision and plans for 2010.</p> <p>Tomas Zabransky Research and Development Manager Center for Addictology Charles University / 1st Medical Faculty / Psychiatric Clinic Czech Republic (2009). Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Reference Group to the UN on HIV and Injecting Drug Use.</p> <p>Naibova, A. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Azerbaijan, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 16 May 2009.</p>		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	3.83	5.21	6.81
Reference	<p>Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.</p> <p>Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.</p>		
General population 15-64 years of age at year of reported data	5 990 102 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	229 500	312 000	407 500
INDICATOR: Percentage of IDUs receiving MMT in a 12 month period	Low	Mid	High
	NK	NK	NK

## OST 25: Number of IDUs receiving BMT in a 12 month period

Data reported	BMT not available		
Year of reported data	2009		
Reference	<p>Naibova, A. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Azerbaijan, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 16 May 2009.</p>		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	3.83	5.21	6.81
Reference	<p>Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.</p> <p>Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.</p>		
General population 15-64 years of age at year of reported data	6 104 384 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	234 000	318 000	415 500
INDICATOR: Percentage of IDUs receiving BMT in a 12 month period	Low	Mid	High
	0	0	0

## OST 26: Number of IDUs receiving other forms of OST in a 12 month period

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Percentage of IDUs receiving other forms of OST at a single point in time	Low	Mid	High

**OST: All documents reviewed**

Afet Naibova Azerbaijan (2009). Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Reference Group to the UN on HIV and Injecting Drug Use.

International Harm Reduction Development Program (2008). Harm reduction developments 2008. New York.

Penitentiary Services, Main Medical Department, et al. (2008). HIV/AIDS Knowledge Attitude and Practice (KAP): "Anonymous Survey among Prisoners and Prison Staff in the Penitentiary System of the Republic of Azerbaijan".

Tomas Zabransky Research and Development Manager Center for Addictology Charles University / 1st Medical Faculty / Psychiatric Clinic Czech Republic (2009). Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Azerbaijan". Reference Group to the UN on HIV and Injecting Drug Use.

United Nations Office on Drugs and Crime Office of the I.R. of Iran and Presidency Drug Control Headquarters (2008). Report - Regional seminar on "Reducing harms of drug use in Middle East, West and Central Asia". Tehran.

UNODC (2008). Milestones: October, 2008 Regional Office for Central Asia. UNODC Regional Office for Central Asia.

UNODC (2009). Status of OST provision and plans for 2010.

WHO and UNAIDS (2008). Progress on implementing the Dublin Declaration on partnership to fight HIV/AIDS in Europe & Central Asia WHO.

**3. Other drug treatment (ODT)****ODT 1 - Detoxification****ODT 1.1: Number of sites providing detoxification**

<b>Is detoxification available?</b>	Yes		
<b>Number of sites - data reported</b>	19 detox sites in prisons		
<b>Year of reported data</b>	2008		
<b>Reference</b>	Penitentiary Services, Main Medical Department, et al. (2008). HIV/AIDS Knowledge Attitude and Practice (KAP): "Anonymous Survey among Prisoners and Prison Staff in the Penitentiary System of the Republic of Azerbaijan".		
<b>Adjustments</b>	Subnational or incomplete data available only		
<b>Final/Adjusted data (to be used as indicator numerator)</b>	NK		
<b>IDU prevalence estimate (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	3.83	5.21	6.81
<b>Reference</b>	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.  Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
<b>General population 15-64 years of age at year of reported data</b>	5 990 102 (source: UN Population Division, 2009)		
<b>Estimated number of IDUs at year of reported data (to be used as indicator denominator)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	229 500	312 000	407 500
<b>INDICATOR: Number of detoxification sites per 1000 IDUs</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	NK	NK	NK

## ODT 1.2: Number of IDUs receiving detoxification

<b>Data reported</b>	Detoxification available but number not known		
<b>Year of reported data</b>	2008		
<b>Reference</b>	Penitentiary Services, Main Medical Department, et al. (2008). HIV/AIDS Knowledge Attitude and Practice (KAP): "Anonymous Survey among Prisoners and Prison Staff in the Penitentiary System of the Republic of Azerbaijan".		
<b>Adjustments</b>			
<b>Final/Adjusted data for 12 month period (to be used as indicator numerator)</b>	NK		
<b>Final/Adjusted data for single point in time (to be used as indicator numerator)</b>	NK		
<b>IDU prevalence estimate (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	3.83	5.21	6.81
<b>Reference</b>	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.  Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
<b>General population 15-64 years of age at year of reported data</b>	5 990 102 (source: UN Population Division, 2009)		
<b>Estimated number of IDUs at year of reported data (to be used as indicator denominator)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	229 500	312 000	407 500
<b>INDICATOR: Percentage of IDUs in detoxification in a 12 month period</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	NK	NK	NK
<b>INDICATOR: Percentage of IDUs in detoxification at a single point in time</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	NK	NK	NK

### ODT 1.3: Number of all clients receiving detoxification in a 12 month period and/or at a single point in time

<b>Data reported</b>	Detoxification available but number not known		
<b>Year of reported data</b>	2008		
<b>Reference</b>	Penitentiary Services, Main Medical Department, et al. (2008). HIV/AIDS Knowledge Attitude and Practice (KAP): "Anonymous Survey among Prisoners and Prison Staff in the Penitentiary System of the Republic of Azerbaijan".		
<b>Adjustments</b>			
<b>Final/Adjusted data for 12 month period (to be used as indicator numerator)</b>	NK		
<b>Final/Adjusted data for single point in time (to be used as indicator numerator)</b>	NK		
<b>IDU prevalence estimate (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	3.83	5.21	6.81
<b>Reference</b>	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.  Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
<b>General population 15-64 years of age at year of reported data</b>	5 990 102 (source: UN Population Division, 2009)		
<b>Estimated number of IDUs at year of reported data (to be used as indicator denominator)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	229 500	312 000	407 500
<b>INDICATOR: Ratio of number of all clients in detoxification in a 12 month period : number of IDUs</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	NK	NK	NK
<b>INDICATOR: Ratio of number of all clients in detoxification at a single point in time : number of IDUs</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	NK	NK	NK

## ODT 2 - Residential rehabilitation

## ODT 2.1: Number of sites providing residential rehabilitation

Is residential rehabilitation present?			
Number of sites - data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Number of residential rehabilitation sites per 1000 IDUs	Low	Mid	High

## ODT 2.2: Number of IDUs in residential rehabilitation

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)			
Final/Adjusted data for single point in time (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Percentage of IDUs in residential rehabilitation in a 12 month period	Low	Mid	High
INDICATOR: Percentage of IDUs in residential rehabilitation at a single point in time	Low	Mid	High

## ODT 2.3: Number of all clients in residential rehabilitation

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)			
Final/Adjusted data for single point in time (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Ratio of number of all drug users in residential rehabilitation in a 12 month period : number of IDUs	Low	Mid	High
INDICATOR: Ratio of number of all drug users in residential rehabilitation at a single point in time : number of IDUs	Low	Mid	High

## ODT 3 - Outpatient psychosocial support, counselling and psychotherapy

## ODT 3.1: Number of sites providing outpatient psychosocial support, counselling or psychotherapy treatment

Is outpatient psychosocial support, counselling or psychotherapy present?	Yes		
Number of sites - data reported	Psychotherapy, counselling or psychosocial support present but number not known		
Year of reported data	2009		
Reference	World Health Organisation (2009). Global health ATLAS data.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	3.83	5.21	6.81
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.  Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
General population 15-64 years of age at year of reported data	6 104 384 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	234 000	318 000	415 500
INDICATOR: Number of outpatient psychosocial support, counselling or psychotherapy sites per 1000 IDUs	Low	Mid	High
	NK	NK	NK

## ODT 3.2: Number of IDUs receiving outpatient psychosocial support, counselling or psychotherapy treatment

Data reported	Psychotherapy, counselling or psychosocial support present but number not known		
Year of reported data	2009		
Reference	World Health Organisation (2009). Global health ATLAS data.		
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)	NK		
Final/Adjusted data for single point in time (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	3.83	5.21	6.81
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.  Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
General population 15-64 years of age at year of reported data	6 104 384 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	234 000	318 000	415 500
INDICATOR: Percentage of IDUs in outpatient psychosocial support, counselling or psychotherapy in a 12 month period	Low	Mid	High
	NK	NK	NK
INDICATOR: Percentage of IDUs receiving outpatient psychosocial support, counselling or psychotherapy at a single point in time	Low	Mid	High
	NK	NK	NK

### ODT 3.3: Number of all clients receiving outpatient psychosocial support, counselling or psychotherapy treatment

<b>Data reported</b>	Psychotherapy, counselling or psychosocial support present but number not known		
<b>Year of reported data</b>	2009		
<b>Reference</b>	World Health Organisation (2009). Global health ATLAS data.		
<b>Adjustments</b>			
<b>Final/Adjusted data for 12 month period (to be used as indicator numerator)</b>	NK		
<b>Final/Adjusted data for single point in time (to be used as indicator numerator)</b>	NK		
<b>IDU prevalence estimate (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	3.83	5.21	6.81
<b>Reference</b>	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.  Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
<b>General population 15-64 years of age at year of reported data</b>	6 104 384 (source: UN Population Division, 2009)		
<b>Estimated number of IDUs at year of reported data (to be used as indicator denominator)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	234 000	318 000	415 500
<b>INDICATOR: Ratio of number of all drug users in outpatient psychosocial support, counselling or psychotherapy in a 12 month period : number of IDUs</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	NK	NK	NK
<b>INDICATOR: Ratio of number of all drug users in outpatient psychosocial support, counselling or psychotherapy at a single point in time : number of IDUs</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	NK	NK	NK

## ODT 4 - Peer-based support groups

## ODT 4.1: Number of peer-based support groups sites

Are Peer-based support groups present?			
Number of sites - data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Number of peer-based support groups sites per 1000 IDUs	Low	Mid	High

## ODT 4.2: Number of IDUs in peer-based support groups

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)			
Final/Adjusted data for single point in time (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Percentage of IDUs in peer-based support groups in a 12 month period	Low	Mid	High
INDICATOR: Percentage of IDUs in peer-based support groups at a single point in time	Low	Mid	High

## ODT 4.3: Number of all clients in peer-based support groups

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)			
Final/Adjusted data for single point in time (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Ratio of number of all drug users in peer-based support groups in a 12 month period : number of IDUs	Low	Mid	High
INDICATOR: Ratio of number of all drug users in peer-based support groups at a single point in time : number of IDUs	Low	Mid	High

## ODT 5 - Drug treatment (treatment type not specified)

## ODT 5.1: Number of drug treatment sites (treatment type not specified)

Type of drug treatment available			
Number of sites - data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Number of drug treatment sites per 1000 IDUs	Low	Mid	High

## ODT 5.2: Number of IDUs in drug treatment (treatment type not specified)

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)			
Final/Adjusted data for single point in time (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Percentage of IDUs in drug treatment not specified/drug treatment combined in a 12 month period	Low	Mid	High
INDICATOR: Percentage of IDUs in drug treatment not specified/drug treatment combined at a single point in time	Low	Mid	High

## ODT 5.3: Number of all clients in drug treatment (treatment type not specified)

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)			
Final/Adjusted data for single point in time (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Ratio of number of all drug users in drug treatment not specified/drug treatment combined in a 12 month period : number of IDUs	Low	Mid	High
INDICATOR: Ratio of number of all drug users in drug treatment not specified/drug treatment combined at a single point in time : number of IDUs	Low	Mid	High

## ODT 6 - Drug treatment in prisons

## ODT 6.1: Number of prison drug treatment sites

Type of drug treatment in prisons?	
Number of sites - data reported	
Year of reported data	
Reference	
Adjustments	
Final/Adjusted data	

## ODT 6.2: Number of IDUs in drug treatment in prisons

Data reported	
Year of reported data	
Reference	
Adjustments	
Final/Adjusted data for 12 month period	
Final/Adjusted data for single point in time	

## ODT 6.3: Number of all clients in drug treatment in prisons

Data reported	
Year of reported data	
Reference	
Adjustments	
Final/Adjusted data for 12 month period	
Final/Adjusted data for single point in time	

## ODT 7 - Compulsory detention as response to drug use

## ODT 7.1: Number of compulsory detention sites for drug use

Number of sites - data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Number of compulsory detention sites for drug use per 1000 IDUs	Low	Mid	High

## ODT 7.2: Number of IDUs in compulsory detention sites for drug use

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)			
Final/Adjusted data for single point in time (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Percentage of IDUs in compulsory detention in closed settings in a 12 month period	Low	Mid	High
INDICATOR: Percentage of IDUs in compulsory detention in closed settings at a single point in time	Low	Mid	High

## ODT 7.3: Number of all drug users in compulsory detention sites for drug use

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)			
Final/Adjusted data for single point in time (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Ratio of all drug users in compulsory detention in closed settings in a 12 month period : number of IDUs	Low	Mid	High
INDICATOR: Ratio of all drug users in compulsory detention in closed settings at a single point in time : number of IDUs	Low	Mid	High

**ODT: All documents reviewed**

Penitentiary Services, Main Medical Department, et al. (2008). HIV/AIDS Knowledge Attitude and Practice (KAP): "Anonymous Survey among Prisoners and Prison Staff in the Penitentiary System of the Republic of Azerbaijan".

World Health Organisation (2009). Global health ATLAS data.

**4. HIV testing and counselling (T&C)****T&C 1: Availability of HIV T&C**

Is T&C available?	
Most recent year of data	
Reference	

**T&C 2: Number of sites providing HIV T&C specifically targeting IDUs**

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	0	0	0
INDICATOR: Number of targeted T&C sites per 1000 IDUs	Low	Mid	High

**T&C 3: Number of sites providing HIV T&C sites to the general population and also accessible to IDUs**

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Number of all T&C sites per 1000 IDUs	Low	Mid	High

**T&C 4: Number of Individual IDUs who were tested for HIV in 12 month period**

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	0	0	0
INDICATOR: Percentage of individual IDUs accessing T&Cs in a 12 month period	Low	Mid	High

**T&C 5: Percentage of IDUs who were tested for HIV in a 12 month period and who know the result**

Year of reported data	
Reference	
Data reported for both male and female IDUs	
Adjustments	
Final/Adjusted data	
Data reported for male IDUs	
Adjustments	
Final/Adjusted data	
Data reported for female IDUs	
Adjustments	
Final/Adjusted data	

**T&C: All documents reviewed**

**No references recorded**

**5. Anti-retro-viral (ARV) treatment****ARV 1: Number of sites providing ARV treatment**

<b>Data reported</b>	1 site		
<b>Year of reported data</b>	2008		
<b>Reference</b>	WHO, UNAIDS., et al. (2009). Towards universal access: Scaling up priority HIV/AIDS interventions in the health sector - progress report 2009. Geneva, WHO.		
<b>Adjustments</b>			
<b>Final/Adjusted data (to be used as indicator numerator)</b>	1		
<b>IDU prevalence estimate (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	3.83	5.21	6.81
<b>Reference</b>	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.  Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
<b>General population 15-64 years of age at year of reported data</b>	5 990 102 (source: UN Population Division, 2009)		
<b>Estimated number of IDUs at year of reported data (to be used as indicator denominator)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	229 500	312 000	407 500
<b>HIV among IDU prevalence estimate (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	2	13	24
<b>Reference</b>	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
<b>Estimated number of HIV positive IDUs at year of reported data (to be used as indicator)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	4590	40 560	97 800
<b>INDICATOR: Number of ARV treatment sites per 1000 IDUs</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	0.00	0.00	0.00
<b>INDICATOR: Number of ARV treatment sites per 1000 IDUs living with HIV</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	0.01	0.02	0.22

## ARV 2: Number of IDUs receiving ARVs at a single point in time

Data reported	Number of IDUs on ART not known		
Year of reported data	2008		
Reference	WHO, UNAIDS., et al. (2009). Towards universal access: Scaling up priority HIV/AIDS interventions in the health sector - progress report 2009. Geneva, WHO.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	3.83	5.21	6.81
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.  Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
General population 15-64 years of age at year of reported data	5 990 102 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	229 500	312 000	407 500
HIV among IDU prevalence estimate (%)	Low	Mid	High
	2	13	24
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
Estimated number of HIV positive IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	4590	40 560	97 800
INDICATOR: Ratio of IDUs on ART: 100 IDUs living with HIV	Low	Mid	High
	NK	NK	NK

## ARV 3: Restrictions on IDUs' access to ARVs

Do restrictions exist which limit IDUs' access to ARV treatment?	
Year of reported data	
Reference	

**ARV: All documents reviewed**

Aceijas, C., E. Oppenheimer, et al. (2006). "Antiretroviral treatment for injecting drug users in developing and transitional countries 1 year before the end of the "Treating 3 million by 2005. Making it happen. The WHO strategy" ("3 by 5")." *Addiction* 101(9): 1246-53.

WHO Europe (2006). Update on HIV/AIDS in the European Region: Ensuring universal access to prevention, treatment, care and support services. Technical briefing document. Copenhagen.

WHO, UNAIDS., et al. (2009). Towards universal access: Scaling up priority HIV/AIDS interventions in the health sector - progress report 2009. Geneva, WHO.

## 6. Condoms

### Condoms 1: Availability of condom programs targeting IDUs

Are condom programs targeting IDUs available?	
Comments	
Year	
Reference	

### Condoms 2: Number of sites distributing condoms to IDUs

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Number of condom distribution sites targeting IDU per 1000 IDUs	Low	Mid	High

### Condoms 3: Number of individual IDUs receiving condoms

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Percentage of IDUs receiving condoms in a 12 month period	Low	Mid	High

## Condoms 4: Number of condoms distributed to IDUs

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Total number of condoms distributed per 12 month period per IDU	Low	Mid	High

**Condoms: All documents reviewed**

Penitentiary Services, Main Medical Department, et al. (2008). HIV/AIDS Knowledge Attitude and Practice (KAP): "Anonymous Survey among Prisoners and Prison Staff in the Penitentiary System of the Republic of Azerbaijan".