

This document was generated by the Reference Group to the United Nations on HIV and Injecting Drug Use.

The data included in this report reflect the results of a systematic search and review of data on a range of indicators reflecting the extent of services provided to people who inject drugs.

These data have been searched, reviewed and extracted according to decision rules developed by the Reference Group.

Some data have been adjusted to facilitate estimates of standard indicators across countries.

Details of the process for searching, grading and including data, along with their interpretation, are detailed in:

Mathers B, Degenhardt L, Ali H, Wiessing L, Hickman M, Mattick R, et al. HIV prevention, treatment and care for people who inject drugs: A systematic review of global, regional and national coverage. *The Lancet* 2010. DOI:10.1016/S0140-6736(10)60232-2.

For more information on the Reference Group, please go to: www.IDURefGroup.Com

Section	Page
Estimated number of people who inject drugs and number of injectors who are living with HIV	4
1. Needle and syringe programs (NSPs)	5
NSP 1: Availability of NSPs	
NSP 2: Number of NSP sites	
NSP 3: Number of pharmacy sites where needles and syringes are available for purchase	
NSP 4: Total number of sites where needles and syringes are available	
NSP 5: Number of individual IDUs in contact with NSPs in a 12 month period	
NSP 6: Number of NSP occasions of service (client contacts) in a 12 month period	
NSP 7: Number of needles and syringes distributed by NSPs in a 12 month period	
2. Opioid substitution programs (OST)	10
OST 1: Availability of opioid substitution treatment (OST)	
OST 2: Availability of methadone maintenance treatment (MMT)	
OST 3: Availability of buprenorphine maintenance treatment (BMT)	
OST 4: Availability of other forms of agonist OST	
OST 5: Dose and duration of MMT	
OST 6: Dose and duration of BMT	
OST 7: Total number of all sites providing OST	
OST 8: Number of sites providing MMT	
OST 9: Number of sites providing BMT	
OST 10: Number of sites providing other forms of agonist OST	
OST 11: Number of all clients in any OST at a single point in time	
OST 12: Number of all clients in MMT at a single point in time	
OST 13: Number of all clients in BMT at a single point in time	
OST 14: Number of all clients in other forms of OST at a single point in time	
OST 15: Number of IDUs in any OST at a single point in time	
OST 16: Number of IDUs in MMT at a single point in time	
OST 17: Number of IDUs in BMT at a single point in time	
OST 18: Number of IDUs in other forms of OST at a single point in time	
OST 19: Number of all clients receiving any OST in a 12 month period	
OST 20: Number of all clients receiving MMT in a 12 month period	
OST 21: Number of all clients receiving BMT in a 12 month period	
OST 22: Number of all clients receiving other forms of OST in a 12 month period	
OST 23: Number of IDUs receiving any OST in a 12 month period	
OST 24: Number of IDUs receiving MMT in a 12 month period	

Section	Page
OST 25: Number of IDUs receiving BMT in a 12 month period	
OST 26: Number of IDUs receiving other forms of OST in a 12 month period	
3. Other Drug Treatment (ODT)	24
<hr/>	
ODT 1: Detoxification	
ODT 1.1: Number of sites providing detoxification	
ODT 1.2: Number of IDUs receiving detoxification	
ODT 1.3: Number of all clients receiving detoxification in a 12 month period and/or at a single point in time	
ODT 2: Residential Rehabilitation	
ODT 2.1: Number of sites providing residential rehabilitation	
ODT 2.2: Number of IDUs in residential rehabilitation	
ODT 2.3: Number of all clients in residential rehabilitation	
ODT 3: Outpatient psychosocial support, counselling and psychotherapy	
ODT 3.1: Number of sites providing outpatient psychosocial support, counselling or psychotherapy	
ODT 3.2: Number of IDUs receiving outpatient psychosocial support, counselling or psychotherapy treatment	
ODT 3.3: Number of all clients receiving outpatient psychosocial support, counselling or psychotherapy treatment	
ODT 4: Peer-based support groups	
ODT 4.1: Number of sites providing peer-based support groups	
ODT 4.2: Number of IDUs in peer-based support groups	
ODT 4.3: Number of all clients in peer-based support groups	
ODT 5: Drug treatment (treatment type not specified)	
ODT 5.1: Number of sites providing unspecified drug treatment	
ODT 5.2: Number of IDUs receiving unspecified drug treatment	
ODT 5.3: Number of all clients receiving unspecified drug treatment	
ODT 6: Drug treatment in prisons	
ODT 6.1: Number of drug treatment sites located in prisons	
ODT 6.2: Number of IDUs receiving drug treatment in prisons	
ODT 6.3: Number of all clients receiving drug treatment in prisons	
ODT 7: Compulsory drug treatment	
ODT 7.1: Number of compulsory detention sites for drug use	
ODT 7.2: Number of IDUs in compulsory detention sites for drug use	
ODT 7.3: Number of all drug users in compulsory detention sites for drug use	
4. HIV testing and counselling (T&C)	38
<hr/>	
T&C 1: Availability of HIV T&C	
T&C 2: Number of sites providing HIV T&C that specifically targets IDUs	

Section	Page
T&C 3: Number of T&C sites providing HIV T& to the general population and also accessible to IDUs	
T&C 4: Number of individual IDUs who were tested for HIV in a 12 month period	
T&C 5: Percentage of IDUs who were tested for HIV in a 12 month period and who know the result	
5. Anti-retro-viral (ARV) treatment	41
<hr/>	
ARV 1: Number of sites providing ARV treatment	
ARV 2: Number of IDUs receiving ARVs at a single point in time	
ARV 3: Restrictions on IDUs' access to ARVs	
6. Condoms	44
<hr/>	
Condoms 1: Availability of condom programs targeting IDUs	
Condoms 2: Number of sites distributing condoms to IDUs	
Condoms 3: Number of individual IDUs receiving condoms	
Condoms 4: Number of condoms distributed to IDUs	

Bangladesh (South Asia)

Estimated number of people who inject drugs and number of injectors who are living with HIV

Estimated prevalence of IDU among general population aged 15-64 years (%)	Low	Mid	High
	0.02	0.03	0.04
Year of IDU prevalence estimate	2005		
Reference for IDU prevalence estimate	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
Estimated prevalence of HIV among people who inject (%)	Low	Mid	High
	0.8	1.35	1.9
Year of HIV prevalence among IDU estimate	2006		
Reference for HIV prevalence among IDU estimate	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age in 2008	102 621 586 (source: UN Population Division, 2009)		
Estimated IDU population in 2008	Low	Mid	High
	22 000	33 000	43 500
Estimated number of IDUs living with HIV in 2008	Low	Mid	High
	176	446	826

1. Needle and syringe programs (NSPs)

NSP 1: Availability of NSPs

Are NSPs available?	Yes
Comments	
Most recent year of documentation	2008
Reference	See references for other NSP indicators
When were NSPs first introduced?	1998
Reference	<p>Lawyers Collective HIV/AIDS Unit (2007). Legal and policy concerns related to IDU harm reduction in SAARC countries. Regional Office for South Asia, United Nations Office on Drugs and Crime</p> <p>Panda, S. and M. Sharma (2006). "Needle syringe acquisition and HIV prevention among injecting drug users: a treatise on the "good" and "not so good" public health practices in South Asia." Substance Use & Misuse 41(6-7): 953-77.</p> <p>Foss, A. M., C. H. Watts, et al. (2007). "Could the CARE-SHAKTI intervention for injecting drug users be maintaining the low HIV prevalence in Dhaka, Bangladesh?" Addiction 102(1): 114-25.</p> <p>Azim, T., N. Hussein, et al. (2005). "Effectiveness of harm reduction programmes for injecting drug users in Dhaka city." Harm Reduction Journal 2(1): 22.</p>
Are needles and syringes available for purchase from pharmacies?	Yes
Most recent year of documentation	2009
Reference	Azim, T. Personal Communication: Response to "Bangladesh - UN Reference Group - Final Data Check". 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 4 October 2009.

NSP 2: Number of NSP sites

Data reported	93 NSPs		
Year of reported data	2008		
Reference	<p>WHO, UNAIDS., et al. (2009). Towards universal access: Scaling up priority HIV/AIDS interventions in the health sector - progress report 2009. Geneva, WHO.</p> <p>Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use</p> <p>UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief on injecting drug use and HIV/AIDS (Draft).</p>		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	93		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Number of NSP sites per 1000 IDUs	Low	Mid	High
	2.14	2.82	4.23

NSP 3: Number of pharmacy sites where needles and syringes are available for purchase

Data reported	Needles and syringes available for purchase from pharmacies but number of sites not available		
Year of reported data	2009		
Reference	Azim, T. Personal Communication: Response to "Bangladesh - UN Reference Group - Final Data Check". 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 4 October 2009.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	2005		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Number of pharmacy sites per 1000 IDUs	Low	Mid	High
	NK	NK	NK

NSP 4: Total number of sites where needles and syringes are available

Total number (NSP sites + Pharmacy sites)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data	(source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Total number of sites per 1000 IDUs	Low	Mid	High

NSP 5: Number of individual IDUs in contact with NSPs in a 12 month period

Data reported	23 684 IDUs accessing services in 2008 32 766 IDUs received syringes between July - Sep 2008		
Year of reported data	2008		
Reference	UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief on injecting drug use and HIV/AIDS (Draft). and Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use Fowler, D. (2008). The HIV/AIDS Prevention Program (HAPP) and HIV/AIDS Targeted Intervention Program (HATI): Documentation and Lessons Learnt in a Program of HIV Targeted Interventions for High Risk Groups in Bangladesh. Draft Report. Dhaka.		
Adjustments	Figures taken as a range		
Final/Adjusted data (to be used as indicator numerator)	23 684 - 32 766		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Percentage of IDUs in contact with NSPs in a 12 month period	Low	Mid	High
	54.4	92.7	148.9

NSP 6: Number of NSP occasions of service (client contacts) in a 12 month period

Data reported	NSP present but number not known		
Year of reported data	2008		
Reference	<p>Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication).</p> <p>UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief on injecting drug use and HIV/AIDS (Draft). and Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use</p> <p>Fowler, D. (2008). The HIV/AIDS Prevention Program (HAPP) and HIV/AIDS Targeted Intervention Program (HATI): Documentation and Lessons Learnt in a Program of HIV Targeted Interventions for High Risk Groups in Bangladesh. Draft Report. Dhaka.</p>		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Number of NSP occasions of service (client contacts) in a 12 month period per IDU	Low	Mid	High
	NK	NK	NK

NSP 7: Number of needles and syringes distributed by NSPs in a 12 month period

Data reported	3 696 224 syringes and needles distributed		
	4 072 729 syringes distributed		
Year of reported data	2008		
Reference	<p>Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use</p> <p>World Health Organization (2009) Universal Access Unpublished Data.</p>		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	3 696 224 - 4 072 729		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Total number of needles/syringes distributed in a 12 month period per IDU	Low	Mid	High
	84.97	117.7	185.12

NSP: All documents reviewed

Asian Harm Reduction Network (AHRN) (2004). Scaling up provision of anti-retrovirals to injecting drug users and non-injecting drug users in Asia: World Health Organization "3 x 5" initiative, Asian Harm Reduction Network (AHRN).

Azim, T. Personal Communication: Response to "Bangladesh - UN Reference Group - Final Data Check". 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 4 October 2009.

Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use

Azim, T., N. Hussein, et al. (2005). "Effectiveness of harm reduction programmes for injecting drug users in Dhaka city." Harm Reduction Journal

Burrows, D. (2006). "Advocacy and coverage of needle exchange programs: results of a comparative study of harm reduction programs in Brazil, Bangladesh, Belarus, Ukraine, Russian Federation, and China." *Cadernos de Saude Publica* 22(4): 871-9.

Burrows, D. (2006). High coverage sites: HIV prevention among injecting drug users in transitional and developing countries: case studies. UNAIDS best practice collection, UNAIDS.

Country coordinating mechanism (2007). Seventh Round Proposal - Country coordination mechanism: Bangladesh. Dhaka: 103.

Foss, A. M., C. H. Watts, et al. (2007). "Could the CARE-SHAKTI intervention for injecting drug users be maintaining the low HIV prevalence in Dhaka, Bangladesh?" *Addiction* 102(1): 114-25.

Fowler, D. (2008). The HIV/AIDS Prevention Program (HAPP) and HIV/AIDS Targeted Intervention Program (HATI): Documentation and Lessons Learnt in a Program of HIV Targeted Interventions for High Risk Groups in Bangladesh. Draft Report. Dhaka.

Hoque, M. M. (HIV/AIDS Advisor), UNODC Bangladesh. Personal Communication: Information for the Reference Group. Bangladesh, 2008. Received by the Reference Group to the UN on HIV and Injecting Drug Use

International Harm Reduction Association (IHRA) (2007). Conference abstracts book. 18th International Conference on the Reduction of Drug Related Harm, Warsaw, Poland.

Islam, M. M. and K. M. Conigrave (2007). "Increasing prevalence of HIV, and persistent high-risk behaviours among drug users in Bangladesh: need for a comprehensive harm reduction programme." *Drug & Alcohol Review* 26(4): 445-54.

Knoess, J. (2006). HIV/AIDS and Intravenous Drug Use in Asia: A mapping of existing HIV/AIDS and harm reduction programs, Development Oriented Drug Control Program GTZ Eschborn: 179.

Knoess, J. (2006). HIV/AIDS and Intravenous Drug Use in Asia: A mapping of existing HIV/AIDS and Harm Reduction Programs, Development Oriented Drug Control Program.

Lawyers Collective HIV/AIDS Unit (2007). Legal and policy concerns related to IDU Harm Reduction in SAARC countries Prevention of transmission of HIV among drug users in SAARC countries. UNODC.

Lawyers Collective HIV/AIDS Unit (2007). Legal and policy concerns related to IDU harm reduction in SAARC countries. Regional Office for South Asia, United Nations Office on Drugs and Crime

Monitoring AIDS Pandemic (MAP) Network (2004). AIDS in Asia: face the facts. A comprehensive analysis of the AIDS epidemics in Asia, Monitoring AIDS Pandemic (MAP) Network.

Monitoring AIDS Pandemic Network (2005). MAP Report 2005: Drug injection and HIV/AIDS in Asia.

National AIDS/STD Programme Bangladesh (2008). 2008 UNGASS Country Progress Report: Bangladesh. Reporting period: January 2006-December 2007. Dhaka, Ministry of Health and Family Welfare: 42.

Needle, R., D. Burrows, et al. (2005). "Effectiveness of community-based outreach in preventing HIV/AIDS among injecting drug users." *International Journal of Drug Policy* 16(Supplement 1): S45-S57.

Panda, S. and M. Sharma (2006). "Needle syringe acquisition and HIV prevention among injecting drug users: a treatise on the "good" and "not so good" public health practices in South Asia." *Substance Use & Misuse* 41(6-7): 953-77.

Reddy, A., M. M. Hoque, et al. (2008). "HIV transmission in Bangladesh: an analysis of IDU programme coverage." *International Journal of Drug Policy* 19 Suppl 1: S37-46.

Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." *AIDS* (Accepted for publication).

The Macfarlane Burnet Institute for Medical Research & Public Health Ltd (2005). Policy and programming guide for HIV/AIDS prevention and care among injecting drug users. Melbourne, World Health Organisation: department of HIV/AIDS.

Todd, C. S., B. Nassiramanesh, et al. (2007). "Emerging HIV epidemics in Muslim countries: assessment of different cultural responses to harm reduction and implications for HIV control." *Current HIV/AIDS Reports* 4(4): 151-7.

UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief on injecting drug use and HIV/AIDS (Draft).

NSP: All documents reviewed

UNAIDS (2006). High coverage sites - HIV prevention among injecting drug users in transitional and developing countries. UNAIDS best practice collection. Geneva.

UNODC. (2007). "State of the HIV Epidemic: Afghanistan, Bangladesh, India, Myanmar, Nepal, Pakistan." from http://www.unodc.org/india/country_profiles.html.

WHO, UNAIDS., et al. (2009). Towards universal access: Scaling up priority HIV/AIDS interventions in the health sector - progress report 2009. Geneva, WHO.

2. Opioid substitution treatment (OST)**OST 1: Availability of opioid substitution treatment (OST)**

Are any forms of agonist OST available?	No
Comments	
Most recent year of documentation	2009
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use
When was agonist OST first available?	
Reference	

OST 2: Availability of methadone maintenance treatment (MMT)

Is MMT available?	No
Comments	
Most recent year of documentation	2009
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use
When was MMT first available?	
Reference	

OST 3: Availability of buprenorphine maintenance treatment (BMT)

Is BMT available?	No
Comments	
Most recent year of documentation	2009
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use
When was BMT first available?	
Reference	

OST 4: Availability of other forms of agonist OST

Are other forms of OST available?	No
Comments	
Most recent year of documentation	2009
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use
When were other forms of OST first available?	
Reference	

OST 5: Dose and duration of MMT

Average/standard maintenance dose of MMT	
Comments	
Year	
Reference	
Average/standard duration of MMT (range)	
Comments	
Year	
Reference	

OST 6: Dose and duration of BMT

Average/standard maintenance dose of BMT	
Comments	
Year	
Reference	
Average/standard duration of BMT (range)	
Comments	
Year	
Reference	

OST 7: Total number of all sites providing OST

Data reported	In 2008 Government gave approval for pilot MMT program for 200 clients in 3 centres in Dhaka but yet to be implemented.		
Year of reported data	2009		
Reference	WHO, UNAIDS., et al. (2009). Towards universal access: Scaling up priority HIV/AIDS interventions in the health sector - progress report 2009. Geneva, WHO. Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Number of OST sites per 1000 IDUs	Low	Mid	High
	0	0	0

OST 8: Number of sites providing MMT

Data reported	In 2008 Government gave approval for pilot MMT program for 200 clients in 3 centres in Dhaka but yet to be implemented.		
Year of reported data	2009		
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Number of MMT sites per 1000 IDUs	Low	Mid	High
	0	0	0

OST 9: Number of sites providing BMT

Data reported	BMT not available		
Year of reported data	2009		
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Number of BMT sites per 1000 IDUs	Low	Mid	High
	0	0	0

OST 10: Number of sites providing other forms of agonist OST

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Number of sites providing other forms of OST per 1000 IDUs	Low	Mid	High

OST 11: Number of all clients in any OST at a single point in time

Data reported	OST not available		
Year of reported data	2009		
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on any OST at census date : 100 IDUs	Low	Mid	High
	0	0	0

OST 12: Number of all clients in MMT at a single point in time

Data reported	OST not available		
Year of reported data	2009		
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on MMT at census date : 100 IDUs	Low	Mid	High
	0	0	0

OST 13: Number of all clients in BMT at a single point in time

Data reported	OST not available		
Year of reported data	2009		
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on BMT at census date : 100 IDUs	Low	Mid	High
	0	0	0

OST 14: Number of all clients in other forms of OST at a single point in time

Data reported	OST not available		
Year of reported data	2009		
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on other forms of OST at census date : 100 IDUs	Low	Mid	High
	0	0	0

OST 15: Number of IDUs in any OST at a single point in time

Data reported	OST not available		
Year of reported data	2009		
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Percentage of IDUs in OST at a single point in time	Low	Mid	High
	0	0	0

OST 16: Number of IDUs in MMT at a single point in time

Data reported	OST not available		
Year of reported data	2009		
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Percentage of IDUs in MMT at a single point in time	Low	Mid	High
	0	0	0

OST 17: Number of IDUs in BMT at a single point in time

Data reported	OST not available		
Year of reported data	2009		
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Percentage of IDUs in BMT at a single point in time	Low	Mid	High
	0	0	0

OST 18: Number of IDUs in other forms of OST at a single point in time

Data reported	OST not available		
Year of reported data	2009		
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Percentage of IDUs in other forms of OST at a single point in time	Low	Mid	High
	0	0	0

OST 19: Number of all clients receiving any OST in a 12 month period

Data reported	OST not available		
Year of reported data	2009		
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on any OST in a 12 month period : 100 IDUs	Low	Mid	High
	0	0	0

OST 20: Number of all clients receiving MMT in a 12 month period

Data reported	OST not available		
Year of reported data	2009		
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on MMT in a 12 month period : 100 IDUs	Low	Mid	High
	0	0	0

OST 21: Number of all clients receiving BMT in a 12 month period

Data reported	OST not available		
Year of reported data	2009		
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on BMT in a 12 month period : 100 IDUs	Low	Mid	High
	0	0	0

OST 22: Number of all clients receiving other forms of OST in a 12 month period

Data reported	OST not available		
Year of reported data	2009		
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on other forms of OST in a 12 month period : 100 IDUs	Low	Mid	High
	0	0	0

OST 23: Number of IDUs receiving any OST in a 12 month period

Data reported	OST not available		
Year of reported data	2009		
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Percentage of IDUs receiving OST in a 12 month period	Low	Mid	High
	0	0	0

OST 24: Number of IDUs receiving MMT in a 12 month period

Data reported	OST not available		
Year of reported data	2009		
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Percentage of IDUs receiving MMT in a 12 month period	Low	Mid	High
	0	0	0

OST 25: Number of IDUs receiving BMT in a 12 month period

Data reported	OST not available		
Year of reported data	2009		
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Percentage of IDUs receiving BMT in a 12 month period	Low	Mid	High
	0	0	0

OST 26: Number of IDUs receiving other forms of OST in a 12 month period

Data reported	OST not available		
Year of reported data	2009		
Reference	Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication). Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	0		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
INDICATOR: Percentage of IDUs receiving other forms of OST at a single point in time	Low	Mid	High
	0	0	0

OST: All documents reviewed

-

Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use

Burrows, D. (2006). High coverage sites: HIV prevention among injecting drug users in transitional and developing countries: case studies. UNAIDS best practice collection, UNAIDS.

Hoque, M. M. (HIV/AIDS Advisor), UNODC Bangladesh. Personal Communication: Information for the Reference Group. Bangladesh, 2008. Received by the Reference Group to the UN on HIV and Injecting Drug Use

Lawyers Collective HIV/AIDS Unit (2007). Legal and policy concerns related to IDU harm reduction in SAARC countries. Regional Office for South Asia, United Nations Office on Drugs and Crime

National AIDS/STD Programme Bangladesh (2008). 2008 UNGASS Country Progress Report: Bangladesh. Reporting period: January 2006-December 2007. Dhaka, Ministry of Health and Family Welfare: 42.

Sharma, M., E. Oppenheimer, et al. (2009). "A situation update on HIV epidemics among people who inject drugs and national responses in South-East Asia Region." AIDS (Accepted for publication).

3. Other drug treatment (ODT)**ODT 1 - Detoxification****ODT 1.1: Number of sites providing detoxification**

Is detoxification available?	yes		
Number of sites - data reported	162 NGO/private clinics provide detoxification 4 centres run by the department of narcotics		
Year of reported data	2008		
Reference	UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief - Injecting drug use and HIV/AIDS.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	166		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 null		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Number of detoxification sites per 1000 IDUs	Low	Mid	High
	3.82	5.03	7.55

ODT 1.2: Number of IDUs receiving detoxification

Data reported	Detoxification available but number not known		
Year of reported data	2008		
Reference	UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief - Injecting drug use and HIV/AIDS.		
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)	NK		
Final/Adjusted data for single point in time (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 null		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Percentage of IDUs in detoxification in a 12 month period	Low	Mid	High
	NK	NK	NK
INDICATOR: Percentage of IDUs in detoxification at a single point in time	Low	Mid	High
	NK	NK	NK

ODT 1.3: Number of all clients receiving detoxification in a 12 month period and/or at a single point in time

Data reported	2583 clients received detoxification in 12 month period		
Year of reported data	2008		
Reference	UN Regional Task Force on Injecting Drug USe and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief - Injecting drug use and HIV/AIDS.		
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)	2583		
Final/Adjusted data for single point in time (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 null		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Ratio of number of all clients in detoxification in a 12 month period : number of IDUs	Low	Mid	High
	0.06	0.08	0.12
INDICATOR: Ratio of number of all clients in detoxification at a single point in time : number of IDUs	Low	Mid	High
	NK	NK	NK

ODT 2 - Residential rehabilitation

ODT 2.1: Number of sites providing residential rehabilitation

Is residential rehabilitation present?	Yes		
Number of sites - data reported	Residential rehabilitation provided by 10 centres provided by FHI Other NGO services provide rehabilitation but number not known		
Year of reported data	2008		
Reference	UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief - Injecting drug use and HIV/AIDS.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	> 10		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 null		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Number of residential rehabilitation sites per 1000 IDUs	Low	Mid	High
	0.23	0.30	0.45

ODT 2.2: Number of IDUs in residential rehabilitation

Data reported	Residential rehabilitation present but number not known		
Year of reported data	2008		
Reference	UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief - Injecting drug use and HIV/AIDS.		
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)	NK		
Final/Adjusted data for single point in time (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 null		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Percentage of IDUs in residential rehabilitation in a 12 month period	Low	Mid	High
	NK	NK	NK
INDICATOR: Percentage of IDUs in residential rehabilitation at a single point in time	Low	Mid	High
	NK	NK	NK

ODT 2.3: Number of all clients in residential rehabilitation

Data reported	1209 clients served by drug treatment centres supported by FHI Number of clients receiving rehabilitation from other NGO operated services not known		
Year of reported data	2008		
Reference	UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief - Injecting drug use and HIV/AIDS.		
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)	> 1209		
Final/Adjusted data for single point in time (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 null		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Ratio of number of all drug users in residential rehabilitation in a 12 month period : number of IDUs	Low	Mid	High
	0.03	0.04	0.05
INDICATOR: Ratio of number of all drug users in residential rehabilitation at a single point in time : number of IDUs	Low	Mid	High
	NK	NK	NK

ODT 3 - Outpatient psychosocial support, counselling and psychotherapy

ODT 3.1: Number of sites providing outpatient psychosocial support, counselling or psychotherapy treatment

Is outpatient psychosocial support, counselling or psychotherapy present?	yes		
Number of sites - data reported	162 NGO/private clinics provide detoxification		
Year of reported data	2008		
Reference	UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief - Injecting drug use and HIV/AIDS.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	162		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 null		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Number of outpatient psychosocial support, counselling or psychotherapy sites per 1000 IDUs	Low	Mid	High
	3.72	4.91	7.36

ODT 3.2: Number of IDUs receiving outpatient psychosocial support, counselling or psychotherapy treatment

Data reported	Psychosocial support and counselling present but number not known		
Year of reported data	2008		
Reference	Hoque, M. M. (HIV/AIDS Advisor), UNODC Bangladesh. Personal Communication: Information for the Reference Group. Bangladesh, 2008. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)	NK		
Final/Adjusted data for single point in time (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 null		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Percentage of IDUs in outpatient psychosocial support, counselling or psychotherapy in a 12 month period	Low	Mid	High
	NK	NK	NK
INDICATOR: Percentage of IDUs receiving outpatient psychosocial support, counselling or psychotherapy at a single point in time	Low	Mid	High
	NK	NK	NK

ODT 3.3: Number of all clients receiving outpatient psychosocial support, counselling or psychotherapy treatment

Data reported	Psychosocial support and counselling present but number not known		
Year of reported data	2008		
Reference	Hoque, M. M. (HIV/AIDS Advisor), UNODC Bangladesh. Personal Communication: Information for the Reference Group. Bangladesh, 2008. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)	NK		
Final/Adjusted data for single point in time (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 null		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Ratio of number of all drug users in outpatient psychosocial support, counselling or psychotherapy in a 12 month period : number of IDUs	Low	Mid	High
	NK	NK	NK
INDICATOR: Ratio of number of all drug users in outpatient psychosocial support, counselling or psychotherapy at a single point in time : number of IDUs	Low	Mid	High
	NK	NK	NK

ODT 4 - Peer-based support groups

ODT 4.1: Number of peer-based support groups sites

Are Peer-based support groups present?	Yes		
Number of sites - data reported	90 NA meetings per week		
Year of reported data	2008		
Reference	Narcotics Anonymous World Services (2009). Quadrennial report for non-governmental organizations - July 2009.		
Adjustments	Assumed each meeting per week occurs at a separate location		
Final/Adjusted data (to be used as indicator numerator)	90		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 null		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Number of peer-based support groups sites per 1000 IDUs	Low	Mid	High
	2.07	2.73	4.09

ODT 4.2: Number of IDUs in peer-based support groups

Data reported	Narcotics Anonymous present but number not known		
Year of reported data	2008		
Reference	Narcotics Anonymous World Services (2009). Quadrennial report for non-governmental organizations - July 2009.		
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)	NK		
Final/Adjusted data for single point in time (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 null		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Percentage of IDUs in peer-based support groups in a 12 month period	Low	Mid	High
	NK	NK	NK
INDICATOR: Percentage of IDUs in peer-based support groups at a single point in time	Low	Mid	High
	NK	NK	NK

ODT 4.3: Number of all clients in peer-based support groups

Data reported	Narcotics Anonymous present but number not known		
Year of reported data	2008		
Reference	Narcotics Anonymous World Services (2009). Quadrennial report for non-governmental organizations - July 2009.		
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)	NK		
Final/Adjusted data for single point in time (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 null		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Ratio of number of all drug users in peer-based support groups in a 12 month period : number of IDUs	Low	Mid	High
	NK	NK	NK
INDICATOR: Ratio of number of all drug users in peer-based support groups at a single point in time : number of IDUs	Low	Mid	High
	NK	null	null

ODT 5 - Drug treatment (treatment type not specified)

ODT 5.1: Number of drug treatment sites (treatment type not specified)

Type of drug treatment available	Unspecified		
Number of sites - data reported	4 centres operated by Department of Narcotics Control, 162 centres by NGOs (includes 10 centres by FHI)		
Year of reported data	2008		
Reference	UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief - Injecting drug use and HIV/AIDS.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	166		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Number of drug treatment sites per 1000 IDUs	Low	Mid	High
	3.82	5.03	7.55

ODT 5.2: Number of IDUs in drug treatment (treatment type not specified)

Data reported	3164 IDUs covered by demand reduction including detoxification		
Year of reported data	2006		
Reference	Reddy, A., M. M. Hoque, et al. (2008). "HIV transmission in Bangladesh: an analysis of IDU programme coverage." International Journal of Drug Policy 19 Suppl 1: S37-46.		
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)	3164		
Final/Adjusted data for single point in time (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	98 058 601 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	21 000	31 500	42 000
INDICATOR: Percentage of IDUs in drug treatment not specified/drug treatment combined in a 12 month period	Low	Mid	High
	7.53	10.04	15.07
INDICATOR: Percentage of IDUs in drug treatment not specified/drug treatment combined at a single point in time	Low	Mid	High
	NK	NK	NK

ODT 5.3: Number of all clients in drug treatment (treatment type not specified)

Data reported	Drug treatment available but number of clients in treatment not known		
Year of reported data	2006		
Reference	Reddy, A., M. M. Hoque, et al. (2008). "HIV transmission in Bangladesh: an analysis of IDU programme coverage." International Journal of Drug Policy 19 Suppl 1: S37-46.		
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)	NK		
Final/Adjusted data for single point in time (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	98 058 601 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	21 000	31 500	42 000
INDICATOR: Ratio of number of all drug users in drug treatment not specified/drug treatment combined in a 12 month period : number of IDUs	Low	Mid	High
	NK	NK	NK
INDICATOR: Ratio of number of all drug users in drug treatment not specified/drug treatment combined at a single point in time : number of IDUs	Low	Mid	High
	NK	NK	NK

ODT 6 - Drug treatment in prisons

ODT 6.1: Number of prison drug treatment sites

Type of drug treatment in prisons?	
Number of sites - data reported	4 drug treatment sites operated by directorate of prisons situated in prison hospitals these sites provide detoxification
Year of reported data	2008
Reference	Hoque, M. M. (HIV/AIDS Advisor), UNODC Bangladesh. Personal Communication: Information for the Reference Group. Bangladesh, 2008. Received by the Reference Group to the UN on HIV and Injecting Drug Use UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief - Injecting drug use and HIV/AIDS.
Adjustments	
Final/Adjusted data	4

ODT 6.2: Number of IDUs in drug treatment in prisons

Data reported	Drug treatment in prisons available but number of IDUs in treatment not known
Year of reported data	2008
Reference	Hoque, M. M. (HIV/AIDS Advisor), UNODC Bangladesh. Personal Communication: Information for the Reference Group. Bangladesh, 2008. Received by the Reference Group to the UN on HIV and Injecting Drug Use UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief - Injecting drug use and HIV/AIDS.
Adjustments	
Final/Adjusted data for 12 month period	NK
Final/Adjusted data for single point in time	NK

ODT 6.3: Number of all clients in drug treatment in prisons

Data reported	2073 clients received treatment at prison drug treatment sites
Year of reported data	2008
Reference	UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief on injecting drug use and HIV/AIDS.
Adjustments	
Final/Adjusted data for 12 month period	2073
Final/Adjusted data for single point in time	NK

ODT 7 - Compulsory detention as response to drug use

ODT 7.1: Number of compulsory detention sites for drug use

Number of sites - data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Number of compulsory detention sites for drug use per 1000 IDUs	Low	Mid	High

ODT 7.2: Number of IDUs in compulsory detention sites for drug use

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)			
Final/Adjusted data for single point in time (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Percentage of IDUs in compulsory detention in closed settings in a 12 month period	Low	Mid	High
INDICATOR: Percentage of IDUs in compulsory detention in closed settings at a single point in time	Low	Mid	High

ODT 7.3: Number of all drug users in compulsory detention sites for drug use

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)			
Final/Adjusted data for single point in time (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Ratio of all drug users in compulsory detention in closed settings in a 12 month period : number of IDUs	Low	Mid	High
INDICATOR: Ratio of all drug users in compulsory detention in closed settings at a single point in time : number of IDUs	Low	Mid	High

ODT: All documents reviewed

Asian Harm Reduction Network (AHRN) (2004). Scaling up provision of anti-retrovirals to injecting drug users and non-injecting drug users in Asia: World Health Organization "3 x 5" initiative, Asian Harm Reduction Network (AHRN).

Azim, T. Personal Communication: Response to "Bangladesh - UN Reference Group - Final Data Check". 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use on 4 October 2009.

Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use

Azim, T., N. Hussein, et al. (2005). "Effectiveness of harm reduction programmes for injecting drug users in Dhaka city." Harm Reduction Journal

Burrows, D. (2006). "Advocacy and coverage of needle exchange programs: results of a comparative study of harm reduction programs in Brazil, Bangladesh, Belarus, Ukraine, Russian Federation, and China." *Cadernos de Saude Publica* 22(4): 871-9.

Burrows, D. (2006). High coverage sites: HIV prevention among injecting drug users in transitional and developing countries: case studies. UNAIDS best practice collection, UNAIDS.

Dhaka Ahsania Mission (2008). Annual Narrative Report on HIV and STI Prevention Project for (I) DUs. Dhaka.

Dhaka Ahsania Mission Report of DAM.

Hoque, M. M. (HIV/AIDS Advisor), UNODC Bangladesh. Personal Communication: Information for the Reference Group. Bangladesh, 2008. Received by the Reference Group to the UN on HIV and Injecting Drug Use

Islam, M. M. and K. M. Conigrave (2007). "Increasing prevalence of HIV, and persistent high-risk behaviours among drug users in Bangladesh: need for a comprehensive harm reduction programme." *Drug & Alcohol Review* 26(4): 445-54.

Knoess, J. (2006). HIV/AIDS and Intravenous Drug Use in Asia: A mapping of existing HIV/AIDS and harm reduction programs, Development Oriented Drug Control Program GTZ Eschborn: 179.

Knoess, J. (2006). HIV/AIDS and Intravenous Drug Use in Asia: A mapping of existing HIV/AIDS and Harm Reduction Programs, Development Oriented Drug Control Program.

Lawyers Collective HIV/AIDS Unit (2007). Legal and policy concerns related to IDU Harm Reduction in SAARC countries Prevention of transmission of HIV among drug users in SAARC countries. UNODC.

Lawyers Collective HIV/AIDS Unit (2007). Legal and policy concerns related to IDU harm reduction in SAARC countries. Regional Office for South Asia, United Nations Office on Drugs and Crime

National AIDS/STD Programme Bangladesh (2008). 2008 UNGASS Country Progress Report: Bangladesh. Reporting period: January 2006-December 2007. Dhaka, Ministry of Health and Family Welfare: 42.

Panda, S. and M. Sharma (2006). "Needle syringe acquisition and HIV prevention among injecting drug users: a treatise on the "good" and "not so good" public health practices in South Asia." *Substance Use & Misuse* 41(6-7): 953-77.

Reddy, A., M. M. Hoque, et al. (2008). "HIV transmission in Bangladesh: an analysis of IDU programme coverage." *International Journal of Drug Policy* 19 Suppl 1: S37-46.

UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief - Injecting drug use and HIV/AIDS.

UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief on injecting drug use and HIV/AIDS (Draft).

UNAIDS (2006). High coverage sites - HIV prevention among injecting drug users in transitional and developing countries. UNAIDS best practice collection. Geneva.

4. HIV testing and counselling (T&C)

T&C 1: Availability of HIV T&C

Is T&C available?	Yes
Most recent year of data	2008
Reference	See references for other VCT indicators

T&C 2: Number of sites providing HIV T&C specifically targeting IDUs

Data reported	9 VCT centres for IDUs		
Year of reported data	2008		
Reference	UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief on injecting drug use and HIV/AIDS.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	9		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Number of targeted T&C sites per 1000 IDUs	Low	Mid	High
	0.21	0.27	0.41

T&C 3: Number of sites providing HIV T&C sites to the general population and also accessible to IDUs

Data reported	91 VCT centres		
Year of reported data	2008		
Reference	Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief on injecting drug use and HIV/AIDS		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	91		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Number of all T&C sites per 1000 IDUs	Low	Mid	High
	2.09	2.76	4.14

T&C 4: Number of Individual IDUs who were tested for HIV in 12 month period

Data reported	1724 IDUs underwent VCT		
Year of reported data	2008		
Reference	Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief on injecting drug use and HIV/AIDS		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	1724		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Percentage of individual IDUs accessing T&Cs in a 12 month period	Low	Mid	High
	3.96	5.22	7.84

T&C 5: Percentage of IDUs who were tested for HIV in a 12 month period and who know the result

Year of reported data	2008
Reference	WHO, UNAIDS., et al. (2009). Towards universal access: Scaling up priority HIV/AIDS interventions in the health sector - progress report 2009. Geneva, WHO.
Data reported for both male and female IDUs	4% IDUs received an HIV test in the last 12 months and know their results (all male sample)
Adjustments	
Final/Adjusted data	4.00%
Data reported for male IDUs	3% male IDUs received an HIV test in the last 12 months and know their results (all male sample)
Adjustments	
Final/Adjusted data	3.00%
Data reported for female IDUs	
Adjustments	
Final/Adjusted data	

T&C: All documents reviewed

Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use

Balanchivadze, N., D. Otiashvili, et al. (2009). Shadow Report on Drug situation 1998-2008. Alternative Georgia.

Bangladesh National AIDS Committee (2005). UNGAS Indicators Country Report Template: 23.

Country Coordinating Mechanism - Bangladesh (2008). Proposal form - rolling continuation channel. Dhaka: 80.

Dhaka Ahsania Mission (2008). Annual Narrative Report on HIV and STI Prevention Project for (I) DUs. Dhaka.

Fowler, D. (2008). The HIV/AIDS Prevention Program (HAPP) and HIV/AIDS Targeted Intervention Program (HATI): Documentation and Lessons Learnt in a Program of HIV Targeted Interventions for High Risk Groups in Bangladesh. Draft Report. Dhaka.

Lawyers Collective HIV/AIDS Unit (2007). Legal and policy concerns related to IDU Harm Reduction in SAARC countries Prevention of transmission of HIV among drug users in SAARC countries. UNODC.

National AIDS/sTD Program Bangladesh (2004). National strategic plan for HIV/AIDS 2004-2010.

National AIDS/STD Program Bangladesh (2004). National Strategic Plan for HIV/AIDS 2004-2010.

UN Regional Task Force on Injecting Drug Use and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief on injecting drug use and HIV/AIDS (Draft).

UNAIDS (2008). Report on the global AIDS epidemic. Geneva, UNAIDS: 362.

WHO, UNAIDS., et al. (2009). Towards universal access: Scaling up priority HIV/AIDS interventions in the health sector - progress report 2009. Geneva, WHO.

5. Anti-retro-viral (ARV) treatment**ARV 1: Number of sites providing ARV treatment**

Data reported	1 site		
	6 sites		
Year of reported data	2008		
	2009		
Reference	Azim, T. (2009). Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Reference Group to the UN on HIV and Injecting Drug Use. WHO, UNAIDS., et al. (2009). Towards universal access: Scaling up priority HIV/AIDS interventions in the health sector - progress report 2009. Geneva, WHO.		
Adjustments	Assumed 2008 for Universal Access data		
	Taken as range		
Final/Adjusted data (to be used as indicator numerator)	1 - 6		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	104 891 220 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 500	33 500	44 500
HIV among IDU prevalence estimate (%)	Low	Mid	High
	0.80	1.35	1.90
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
Estimated number of HIV positive IDUs at year of reported data (to be used as indicator)	Low	Mid	High
	180	452	846
INDICATOR: Number of ARV treatment sites per 1000 IDUs	Low	Mid	High
	0.02	0.10	0.27
INDICATOR: Number of ARV treatment sites per 1000 IDUs living with HIV	Low	Mid	High
	1.18	7.74	33.33

ARV 2: Number of IDUs receiving ARVs at a single point in time

Data reported	5 IDUs on ART [Defintion of IDUs not given]		
Year of reported data	2008		
Reference	UN Regional Task Force on Injecting Drug USe and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief on injecting drug use and HIV/AIDS. Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	5		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
HIV among IDU prevalence estimate (%)	Low	Mid	High
	0.80	1.35	1.90
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
Estimated number of HIV positive IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	176	446	826
INDICATOR: Ratio of IDUs on ART: 100 IDUs living with HIV	Low	Mid	High
	0.60	1.12	2.84

ARV 3: Restrictions on IDUs' access to ARVs

Do restrictions exist which limit IDUs' access to ARV treatment?	
Year of reported data	
Reference	

ARV: All documents reviewed

Aceijas, C., E. Oppenheimer, et al. (2006). "Antiretroviral treatment for injecting drug users in developing and transitional countries 1 year before the end of the "Treating 3 million by 2005. Making it happen. The WHO strategy" ("3 by 5")." *Addiction* 101(9): 1246-53.

Asian Harm Reduction Network (AHRN) (2004). Scaling up provision of anti-retrovirals to injecting rug users and non-injecting drug users in Asia: World Health Organization "3 x 5" initiative, Asian Harm Reduction Network (AHRN).

Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use

Country coordinating mechanism (2007). Seventh Round Proposal - Country coordination mechanism: Bangladesh. Dhaka: 103.

Country Coordinating Mechanism - Bangladesh (2008). Proposal form - rolling continuation channel. Dhaka: 80.

Lawyers Collective HIV/AIDS Unit (2007). Legal and policy concerns related to IDU harm reduction in SAARC countries. Regional Office for South Asia, United Nations Office on Drugs and Crime

UN Regional Task Force on Injecting Drug USe and HIV/AIDS for Asia and the Pacific (2009). Bangladesh country advocacy brief on injecting drug use and HIV/AIDS (Draft).

WHO, UNAIDS., et al. (2009). Towards universal access: Scaling up priority HIV/AIDS interventions in the health sector - progress report 2009. Geneva, WHO.

6. Condoms

Condoms 1: Availability of condom programs targeting IDUs

Are condom programs targeting IDUs available?	Yes
Comments	
Year	2008
Reference	See references for other Condom indicators

Condoms 2: Number of sites distributing condoms to IDUs

Data reported	Condoms available from NSP sites 93 NSP sites		
Year of reported data	2008		
Reference	Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments	Assumed all NSPs provide condoms		
Final/Adjusted data (to be used as indicator numerator)	93		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." <i>The Lancet</i> 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Number of condom distribution sites targeting IDU per 1000 IDUs	Low	Mid	High
	2.14	2.82	4.23

Condoms 3: Number of individual IDUs receiving condoms

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Percentage of IDUs receiving condoms in a 12 month period	Low	Mid	High

Condoms 4: Number of condoms distributed to IDUs

Data reported	2 775 950 condoms distributed		
Year of reported data	2008		
Reference	Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	2 775 950		
IDU prevalence estimate (%)	Low	Mid	High
	0.02	0.03	0.04
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	102 621 586 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	22 000	33 000	43 500
INDICATOR: Total number of condoms distributed per 12 month period per IDU	Low	Mid	High
	63.81	84.12	126.18

Condoms: All documents reviewed

Azim, T. Personal Communication: Response to the "UN HIV and IDU Reference Group: Request to fill data gaps - Bangladesh". Bangladesh, 2009. Received by the Reference Group to the UN on HIV and Injecting Drug Use

Azim, T., N. Hussein, et al. (2005). "Effectiveness of harm reduction programmes for injecting drug users in Dhaka city." *Harm Reduction Journal*

Burrows, D. (2006). "Advocacy and coverage of needle exchange programs: results of a comparative study of harm reduction programs in Brazil, Bangladesh, Belarus, Ukraine, Russian Federation, and China." *Cadernos de Saude Publica* 22(4): 871-9.

Burrows, D. (2006). High coverage sites: HIV prevention among injecting drug users in transitional and developing countries: case studies. UNAIDS best practice collection, UNAIDS.

Country coordinating mechanism (2007). Seventh Round Proposal - Country coordination mechanism: Bangladesh. Dhaka: 103.

Dhaka Ahsania Mission (2008). Annual Narrative Report on HIV and STI Prevention Project for (I) DUs. Dhaka.

Family Health International (2007). Bangladesh final report: September 1997 - September 2007. Arlington VA: 68.

Foss, A. M., C. H. Watts, et al. (2007). "Could the CARE-SHAKTI intervention for injecting drug users be maintaining the low HIV prevalence in Dhaka, Bangladesh?" *Addiction* 102(1): 114-25.

Fowler, D. (2008). The HIV/AIDS Prevention Program (HAPP) and HIV/AIDS Targeted Intervention Program (HATI): Documentation and Lessons Learnt in a Program of HIV Targeted Interventions for High Risk Groups in Bangladesh. Draft Report. Dhaka.

Needle, R., D. Burrows, et al. (2005). "Effectiveness of community-based outreach in preventing HIV/AIDS among injecting drug users." *International Journal of Drug Policy* 16(Supplement 1): S45-S57.

UNAIDS (2006). High coverage sites - HIV prevention among injecting drug users in transitional and developing countries. UNAIDS best practice collection. Geneva.

UNODC (2008). Numbers at a glance - H13 intervention project.