

**This document was generated by the Reference Group to the United Nations on HIV and Injecting Drug Use.**

**The data included in this report reflect the results of a systematic search and review of data on a range of indicators reflecting the extent of services provided to people who inject drugs.**

**These data have been searched, reviewed and extracted according to decision rules developed by the Reference Group.**

**Some data have been adjusted to facilitate estimates of standard indicators across countries.**

**Details of the process for searching, grading and including data, along with their interpretation, are detailed in:**

Mathers B, Degenhardt L, Ali H, Wiessing L, Hickman M, Mattick R, et al. HIV prevention, treatment and care for people who inject drugs: A systematic review of global, regional and national coverage. *The Lancet* 2010. DOI:10.1016/S0140-6736(10)60232-2.

**For more information on the Reference Group, please go to: [www.IDURefGroup.Com](http://www.IDURefGroup.Com)**

Section	Page
Estimated number of people who inject drugs and number of injectors who are living with HIV	4
<b>1. Needle and syringe programs (NSPs)</b>	<b>5</b>
NSP 1: Availability of NSPs	
NSP 2: Number of NSP sites	
NSP 3: Number of pharmacy sites where needles and syringes are available for purchase	
NSP 4: Total number of sites where needles and syringes are available	
NSP 5: Number of individual IDUs in contact with NSPs in a 12 month period	
NSP 6: Number of NSP occasions of service (client contacts) in a 12 month period	
NSP 7: Number of needles and syringes distributed by NSPs in a 12 month period	
<b>2. Opioid substitution programs (OST)</b>	<b>10</b>
OST 1: Availability of opioid substitution treatment (OST)	
OST 2: Availability of methadone maintenance treatment (MMT)	
OST 3: Availability of buprenorphine maintenance treatment (BMT)	
OST 4: Availability of other forms of agonist OST	
OST 5: Dose and duration of MMT	
OST 6: Dose and duration of BMT	
OST 7: Total number of all sites providing OST	
OST 8: Number of sites providing MMT	
OST 9: Number of sites providing BMT	
OST 10: Number of sites providing other forms of agonist OST	
OST 11: Number of all clients in any OST at a single point in time	
OST 12: Number of all clients in MMT at a single point in time	
OST 13: Number of all clients in BMT at a single point in time	
OST 14: Number of all clients in other forms of OST at a single point in time	
OST 15: Number of IDUs in any OST at a single point in time	
OST 16: Number of IDUs in MMT at a single point in time	
OST 17: Number of IDUs in BMT at a single point in time	
OST 18: Number of IDUs in other forms of OST at a single point in time	
OST 19: Number of all clients receiving any OST in a 12 month period	
OST 20: Number of all clients receiving MMT in a 12 month period	
OST 21: Number of all clients receiving BMT in a 12 month period	
OST 22: Number of all clients receiving other forms of OST in a 12 month period	
OST 23: Number of IDUs receiving any OST in a 12 month period	
OST 24: Number of IDUs receiving MMT in a 12 month period	

<b>Section</b>	<b>Page</b>
OST 25: Number of IDUs receiving BMT in a 12 month period	
OST 26: Number of IDUs receiving other forms of OST in a 12 month period	
<b>3. Other Drug Treatment (ODT)</b>	<b>24</b>
<hr/>	
ODT 1: Detoxification	
ODT 1.1: Number of sites providing detoxification	
ODT 1.2: Number of IDUs receiving detoxification	
ODT 1.3: Number of all clients receiving detoxification in a 12 month period and/or at a single point in time	
ODT 2: Residential Rehabilitation	
ODT 2.1: Number of sites providing residential rehabilitation	
ODT 2.2: Number of IDUs in residential rehabilitation	
ODT 2.3: Number of all clients in residential rehabilitation	
ODT 3: Outpatient psychosocial support, counselling and psychotherapy	
ODT 3.1: Number of sites providing outpatient psychosocial support, counselling or psychotherapy	
ODT 3.2: Number of IDUs receiving outpatient psychosocial support, counselling or psychotherapy treatment	
ODT 3.3: Number of all clients receiving outpatient psychosocial support, counselling or psychotherapy treatment	
ODT 4: Peer-based support groups	
ODT 4.1: Number of sites providing peer-based support groups	
ODT 4.2: Number of IDUs in peer-based support groups	
ODT 4.3: Number of all clients in peer-based support groups	
ODT 5: Drug treatment (treatment type not specified)	
ODT 5.1: Number of sites providing unspecified drug treatment	
ODT 5.2: Number of IDUs receiving unspecified drug treatment	
ODT 5.3: Number of all clients receiving unspecified drug treatment	
ODT 6: Drug treatment in prisons	
ODT 6.1: Number of drug treatment sites located in prisons	
ODT 6.2: Number of IDUs receiving drug treatment in prisons	
ODT 6.3: Number of all clients receiving drug treatment in prisons	
ODT 7: Compulsory drug treatment	
ODT 7.1: Number of compulsory detention sites for drug use	
ODT 7.2: Number of IDUs in compulsory detention sites for drug use	
ODT 7.3: Number of all drug users in compulsory detention sites for drug use	
<b>4. HIV testing and counselling (T&amp;C)</b>	<b>39</b>
<hr/>	
T&C 1: Availability of HIV T&C	
T&C 2: Number of sites providing HIV T&C that specifically targets IDUs	

Section	Page
T&C 3: Number of T&C sites providing HIV T& to the general population and also accessible to IDUs	
T&C 4: Number of individual IDUs who were tested for HIV in a 12 month period	
T&C 5: Percentage of IDUs who were tested for HIV in a 12 month period and who know the result	
<b>5. Anti-retro-viral (ARV) treatment</b>	<b>42</b>
<hr/>	
ARV 1: Number of sites providing ARV treatment	
ARV 2: Number of IDUs receiving ARVs at a single point in time	
ARV 3: Restrictions on IDUs' access to ARVs	
<b>6. Condoms</b>	<b>45</b>
<hr/>	
Condoms 1: Availability of condom programs targeting IDUs	
Condoms 2: Number of sites distributing condoms to IDUs	
Condoms 3: Number of individual IDUs receiving condoms	
Condoms 4: Number of condoms distributed to IDUs	

## Finland (Western Europe)

### Estimated number of people who inject drugs and number of injectors who are living with HIV

Estimated prevalence of IDU among general population aged 15-64 years (%)	Low	Mid	High
	0.35	0.45	0.57
Year of IDU prevalence estimate	2002		
Reference for IDU prevalence estimate	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
Estimated prevalence of HIV among people who inject (%)	Low	Mid	High
	0.10	0.2	0.27
Year of HIV prevalence among IDU estimate	2006		
Reference for HIV prevalence among IDU estimate	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.  Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
General population 15-64 years of age in 2008	3 535 039 (source: UN Population Division, 2009)		
Estimated IDU population in 2008	Low	Mid	High
	12 500	16 000	20 000
Estimated number of IDUs living with HIV in 2008	Low	Mid	High
	13	32	54

## 1. Needle and syringe programs (NSPs)

### NSP 1: Availability of NSPs

Are NSPs available?	Yes
Comments	
Most recent year of documentation	2008
Reference	See references for other NSP indicators
When were NSPs first introduced?	1997
Reference	Finnish National Focal Point - STAKES (2008). Finland Drug Situation - New Development, Trends and in-depth information on selected issues. 2008 National report (2007 data) to the EMCDDA. Helsinki.
Are needles and syringes available for purchase from pharmacies?	
Most recent year of documentation	
Reference	

### NSP 2: Number of NSP sites

Data reported	45 fixed sites 5 outreach workers sites 2 mobile sites		
Year of reported data	2007		
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .		
Adjustments	Assumed counts of all site are mutually exclusive. Total number Of sites calculated.		
Final/Adjusted data (to be used as indicator numerator)	52		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 521 948 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Number of NSP sites per 1000 IDUs	Low	Mid	High
	2.60	3.25	4.16

## NSP 3: Number of pharmacy sites where needles and syringes are available for purchase

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Number of pharmacy sites per 1000 IDUs	Low	Mid	High

## NSP 4: Total number of sites where needles and syringes are available

Total number (NSP sites + Pharmacy sites)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Total number of sites per 1000 IDUs	Low	Mid	High

## NSP 5: Number of individual IDUs in contact with NSPs in a 12 month period

Data reported	13000 individual clients		
Year of reported data	2007		
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .  Reitox National Focal Point (2008). 2008 National Report (2007 data) to the EMCDDA - New developments, trends and in-depth information on selected issues EMCDDA.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	13 000		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 521 948 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Percentage of IDUs in contact with NSPs in a 12 month period	Low	Mid	High
	65	81.2	104

## NSP 6: Number of NSP occasions of service (client contacts) in a 12 month period

Data reported	92000 client contacts		
Year of reported data	2007		
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	92 000		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 521 948 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Number of NSP occasions of service (client contacts) in a 12 month period per IDU	Low	Mid	High
	4.60	5.75	7.36

## NSP 7: Number of needles and syringes distributed by NSPs in a 12 month period

<b>Data reported</b>	264 8000 syringes provided through needle and syringe programmes		
<b>Year of reported data</b>	2007		
<b>Reference</b>	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .		
<b>Adjustments</b>			
<b>Final/Adjusted data (to be used as indicator numerator)</b>	2 648 000		
<b>IDU prevalence estimate (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	0.35	0.45	0.57
<b>Reference</b>	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
<b>General population 15-64 years of age at year of reported data</b>	3 521 948 (source: UN Population Division, 2009)		
<b>Estimated number of IDUs at year of reported data (to be used as indicator denominator)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	12 500	16 000	20 000
<b>INDICATOR: Total number of needles/syringes distributed in a 12 month period per IDU</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	132.40	165.5	211.84

**NSP: All documents reviewed**

EMCDDA (2007). The State of the Drugs Problem in Europe. Luxembourg: 100.

European Monitoring Centre for Drugs and Drug Addiction. (2008). "Health and social responses - Statistical bulletin." Statistical Bulletin Retrieved 6 April 2009, from <http://www.emcdda.europa.eu/stats08/hsr>.

European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <http://www.emcdda.europa.eu/stats09/hsr>.

Finnish National Focal Point - STAKES (2008). Finland Drug Situation - New Development, Trends and in-depth information on selected issues. 2008 National report (2007 data) to the EMCDDA. Helsinki.

Helsinki Deaconess Institute (2008). Organization's structure & work

Ronka, S., Virtanen, A., Perala, R. and Vihmo, J. (2007). Finland Drug situation 2007: New developments, trends and in-depth information on selected issues. Helsinki, Finnish National Focal Point, STAKES: 137.

Salminen, M. (2008). UNGASS HIV/AIDS country progress report Finland. Helsinki, KTL - National Public Health Institute, Finland. Department of Infectious Disease Epidemiology and Control. HIV- Unit.

Salminen, M. (2008). UNGASS HIV/AIDS country progress report Finland. Helsinki, KTL - National Public Health Institute, Finland. Department of Infectious Disease Epidemiology and Control. HIV- Unit.

The Helsinki Deaconess Institutes (2008). Information on HIV services for injecting drug users.

WHO and UNAIDS (2008). Progress on implementing the Dublin Declaration on partnership to fight HIV/AIDS in Europe & Central Asia WHO.

**2. Opioid substitution treatment (OST)****OST 1: Availability of opioid substitution treatment (OST)**

Are any forms of agonist OST available?	Yes
Comments	
Most recent year of documentation	2008
Reference	See references for other OST indicators
When was agonist OST first available?	1974
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .

**OST 2: Availability of methadone maintenance treatment (MMT)**

Is MMT available?	Yes
Comments	
Most recent year of documentation	2008
Reference	See references for other OST indicators
When was MMT first available?	1974
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .

**OST 3: Availability of buprenorphine maintenance treatment (BMT)**

Is BMT available?	Yes
Comments	
Most recent year of documentation	2008
Reference	See references for other OST indicators
When was BMT first available?	1997
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .

**OST 4: Availability of other forms of agonist OST**

Are other forms of OST available?	
Comments	
Most recent year of documentation	
Reference	
When were other forms of OST first available?	
Reference	

## OST 5: Dose and duration of MMT

Average/standard maintenance dose of MMT	60 -120 mg
Comments	
Year	2009
Reference	World Health Organisation (2009). Global health ATLAS data.
Average/standard duration of MMT (range)	
Comments	
Year	
Reference	

## OST 6: Dose and duration of BMT

Average/standard maintenance dose of BMT	16 - 24 mg
Comments	
Year	2009
Reference	World Health Organisation (2009). Global health ATLAS data.
Average/standard duration of BMT (range)	
Comments	
Year	
Reference	

## OST 7: Total number of all sites providing OST

Data reported	OST available but number of sites not known		
Year of reported data	2007		
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 521 948 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Number of OST sites per 1000 IDUs	Low	Mid	High
	NK	NK	NK

## OST 8: Number of sites providing MMT

Data reported	MMT available but number of sites not known		
Year of reported data	2007		
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 521 948 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Number of MMT sites per 1000 IDUs	Low	Mid	High
	NK	NK	NK

## OST 9: Number of sites providing BMT

Data reported	BMT available but number not known		
Year of reported data	2007		
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 521 948 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Number of BMT sites per 1000 IDUs	Low	Mid	High
	NK	NK	NK

## OST 10: Number of sites providing other forms of agonist OST

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Number of sites providing other forms of OST per 1000 IDUs	Low	Mid	High

## OST 11: Number of all clients in any OST at a single point in time

<b>Data reported</b>	500 - 540 clients in MMT  660 clients in BMT [No information on whether number of clients in OST at single point in time (i.e. Census date) or total number of clients receiving OST over a period of time]		
<b>Year of reported data</b>	2009  2007		
<b>Reference</b>	World Health Organisation (2009). Global health ATLAS data.  European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .		
<b>Adjustments</b>	Assumed enrolment at a census date		
<b>Final/Adjusted data (to be used as indicator numerator)</b>	1160 - 1200		
<b>IDU prevalence estimate (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	0.35	0.45	0.57
<b>Reference</b>	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
<b>General population 15-64 years of age at year of reported data</b>	3 542 832 (source: UN Population Division, 2009)		
<b>Estimated number of IDUs at year of reported data (to be used as indicator denominator)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	12 500	16 000	20 000
<b>INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on any OST at census date : 100 IDUs</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	5.80	7.38	9.60

## OST 12: Number of all clients in MMT at a single point in time

<b>Data reported</b>	500 patients on MMT  540 clients in MMT [No information on whether number of clients in MMT at single point in time (i.e. Census date) or total number of clients receiving MMT over a period of time]		
<b>Year of reported data</b>	2009  2007		
<b>Reference</b>	World Health Organisation (2009). Global health ATLAS data.  European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .		
<b>Adjustments</b>	Assumed count equals enrolment at a census date  Figures taken as range		
<b>Final/Adjusted data (to be used as indicator numerator)</b>	500 - 540		
<b>IDU prevalence estimate (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	0.35	0.45	0.57
<b>Reference</b>	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
<b>General population 15-64 years of age at year of reported data</b>	3 542 832 (source: UN Population Division, 2009)		
<b>Estimated number of IDUs at year of reported data (to be used as indicator denominator)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	12 500	16 000	20 000
<b>INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on MMT at census date : 100 IDUs</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	2.50	3.25	4.32

## OST 13: Number of all clients in BMT at a single point in time

Data reported	660 clients in BMT [No information on whether number of clients in BMT at single point in time (i.e. Census date) or total number of clients receiving BMT over a period of time]		
Year of reported data	2007 2009		
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .  World Health Organisation (2009). Global health ATLAS data.		
Adjustments	Assumed count equals enrolment at a census date		
Final/Adjusted data (to be used as indicator numerator)	660		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 542 832 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on BMT at census date : 100 IDUs	Low	Mid	High
	3.30	4.12	5.28

## OST 14: Number of all clients in other forms of OST at a single point in time

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on other forms of OST at census date : 100 IDUs	Low	Mid	High

## OST 15: Number of IDUs in any OST at a single point in time

Data reported	OST available but number of IDU on OST not known		
Year of reported data	2007		
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 521 948 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Percentage of IDUs in OST at a single point in time	Low	Mid	High
	NK	NK	NK

## OST 16: Number of IDUs in MMT at a single point in time

Data reported	MMT available but number of IDUs not known		
Year of reported data	2007		
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 521 948 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Percentage of IDUs in MMT at a single point in time	Low	Mid	High
	NK	NK	NK

## OST 17: Number of IDUs in BMT at a single point in time

Data reported	BMT available but number of IDUs receiving BMT not known		
Year of reported data	2007		
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 521 948 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Percentage of IDUs in BMT at a single point in time	Low	Mid	High
	NK	NK	NK

## OST 18: Number of IDUs in other forms of OST at a single point in time

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Percentage of IDUs in other forms of OST at a single point in time	Low	Mid	High

## OST 19: Number of all clients receiving any OST in a 12 month period

Data reported	540 clients in MMT  660 clients in BMT [No information on whether number of clients in OST at single point in time (i.e. Census date) or total number of clients receiving OST over a period of time]		
Year of reported data	2007		
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .		
Adjustments	Assumed count equal to total number of clients enrolled in OST throughout 2007		
Final/Adjusted data (to be used as indicator numerator)	1200		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 521 948 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on any OST in a 12 month period : 100 IDUs	Low	Mid	High
	6	7.50	9.60

## OST 20: Number of all clients receiving MMT in a 12 month period

Data reported	540 clients in MMT [No information on whether number of clients in MMT at single point in time (i.e. Census date) or total number of clients receiving MMT over a period of time]		
Year of reported data	2007		
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .		
Adjustments	Assumed count equals total number of clients enrolled in MMT throughout 2007		
Final/Adjusted data (to be used as indicator numerator)	540		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 521 948 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on MMT in a 12 month period : 100 IDUs	Low	Mid	High
	2.70	3.38	4.32

## OST 21: Number of all clients receiving BMT in a 12 month period

Data reported	660 clients in BMT [No information on whether number of clients in BMT at single point in time (i.e. Census date) or total number of clients receiving BMT over a period of time]		
Year of reported data	2007		
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .		
Adjustments	Assumed count equals total number of clients enrolled in BMT throughout 2007		
Final/Adjusted data (to be used as indicator numerator)	660		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 521 948 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on BMT in a 12 month period : 100 IDUs	Low	Mid	High
	3.30	4.12	5.28

## OST 22: Number of all clients receiving other forms of OST in a 12 month period

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Ratio of total number of all clients (including IDUs and non-IDUs) on other forms of OST in a 12 month period : 100 IDUs	Low	Mid	High

## OST 23: Number of IDUs receiving any OST in a 12 month period

Data reported	OST available but number of IDU on OST not known		
Year of reported data	2007		
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 521 948 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Percentage of IDUs receiving OST in a 12 month period	Low	Mid	High
	NK	NK	NK

## OST 24: Number of IDUs receiving MMT in a 12 month period

Data reported	MMT available but number of IDUs not known		
Year of reported data	2007		
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 521 948 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Percentage of IDUs receiving MMT in a 12 month period	Low	Mid	High
	NK	NK	NK

## OST 25: Number of IDUs receiving BMT in a 12 month period

Data reported	BMT available but number of IDUs receiving BMT not known		
Year of reported data	2007		
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/hsr">http://www.emcdda.europa.eu/stats09/hsr</a> .		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 521 948 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Percentage of IDUs receiving BMT in a 12 month period	Low	Mid	High
	NK	NK	NK

## OST 26: Number of IDUs receiving other forms of OST in a 12 month period

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Percentage of IDUs receiving other forms of OST at a single point in time	Low	Mid	High

**OST: All documents reviewed**

BISDRO University of Bremen and WIAD Scientific Institute of the German Medical Association Bonn (2008). Reduction of Drug-related Crime in Prison: The impact of opioid substitution treatment on the manageability of opioid dependent prisoners.

Bremen Institute for Drug Research (BISDRO) University of Bremen, Centre for Interdisciplinary Addiction Research (CIAR) University of Hamburg, et al. (2008). Final report on prevention, treatment, and harm reduction services in prison, on reintegration services on release from prison and methods to monitor/analyse drug use among prisoners.

Donoghoe, M. C., A. R. Bollerup, et al. (2007). "Access to highly active antiretroviral therapy (HAART) for injecting drug users in the WHO European Region 2002-2004." *International Journal of Drug Policy* 18(4): 271-80.

European Monitoring Centre for Drugs and Drug Addiction. (2008). "Health and social responses - Statistical bulletin." *Statistical Bulletin* Retrieved 6 April 2009, from <http://www.emcdda.europa.eu/stats08/hsr>.

European Monitoring Centre for Drugs and Drug Addiction. (2009). "Health and social responses - Statistical bulletin." *Statistical Bulletin* 2009, from <http://www.emcdda.europa.eu/stats09/hsr>.

Finnish National Focal Point - STAKES (2008). Finland Drug Situation - New Development, Trends and in-depth information on selected issues. 2008 National report (2007 data) to the EMCDDA. Helsinki.

Ronka, S., Virtanen, A., Perala, R. and Vihmo, J. (2007). Finland Drug situation 2007: New developments, trends and in-depth information on selected issues. Helsinki, Finnish National Focal Point, STAKES: 137.

Salminen, M. (2008). UNGASS HIV/AIDS country progress report Finland. Helsinki, KTL - National Public Health Institute, Finland. Department of Infectious Disease Epidemiology and Control. HIV- Unit.

Solberg, U., G. Burkhart, et al. (2002). "An overview of opiate substitution treatment in the European Union and Norway." *International Journal of Drug Policy* 13(6): 477-484.

The Helsinki Deaconess Institutes (2008). Information on HIV services for injecting drug users.

**3. Other drug treatment (ODT)****ODT 1 - Detoxification****ODT 1.1: Number of sites providing detoxification**

Is detoxification available?	Yes		
Number of sites - data reported	Detoxification available but number not known		
Year of reported data	2009		
Reference	World Health Organisation (2009). Global health ATLAS data.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 542 832 null		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Number of detoxification sites per 1000 IDUs	Low	Mid	High
	NK	NK	NK

**ODT 1.2: Number of IDUs receiving detoxification**

Data reported	Detoxification available but number not known		
Year of reported data	2009		
Reference	World Health Organisation (2009). Global health ATLAS data.		
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)	NK		
Final/Adjusted data for single point in time (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 542 832 null		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Percentage of IDUs in detoxification in a 12 month period	Low	Mid	High
	NK	NK	NK
INDICATOR: Percentage of IDUs in detoxification at a single point in time	Low	Mid	High
	NK	NK	NK

### ODT 1.3: Number of all clients receiving detoxification in a 12 month period and/or at a single point in time

<b>Data reported</b>	Detoxification available but number not known		
<b>Year of reported data</b>	2009		
<b>Reference</b>	World Health Organisation (2009). Global health ATLAS data.		
<b>Adjustments</b>			
<b>Final/Adjusted data for 12 month period (to be used as indicator numerator)</b>	NK		
<b>Final/Adjusted data for single point in time (to be used as indicator numerator)</b>	NK		
<b>IDU prevalence estimate (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	0.35	0.45	0.57
<b>Reference</b>	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
<b>General population 15-64 years of age at year of reported data</b>	3 542 832 null		
<b>Estimated number of IDUs at year of reported data (to be used as indicator denominator)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	12 500	16 000	20 000
<b>INDICATOR: Ratio of number of all clients in detoxification in a 12 month period : number of IDUs</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	NK	NK	NK
<b>INDICATOR: Ratio of number of all clients in detoxification at a single point in time : number of IDUs</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	NK	NK	NK

## ODT 2 - Residential rehabilitation

## ODT 2.1: Number of sites providing residential rehabilitation

Is residential rehabilitation present?			
Number of sites - data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Number of residential rehabilitation sites per 1000 IDUs	Low	Mid	High

## ODT 2.2: Number of IDUs in residential rehabilitation

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)			
Final/Adjusted data for single point in time (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Percentage of IDUs in residential rehabilitation in a 12 month period	Low	Mid	High
INDICATOR: Percentage of IDUs in residential rehabilitation at a single point in time	Low	Mid	High

## ODT 2.3: Number of all clients in residential rehabilitation

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)			
Final/Adjusted data for single point in time (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Ratio of number of all drug users in residential rehabilitation in a 12 month period : number of IDUs	Low	Mid	High
INDICATOR: Ratio of number of all drug users in residential rehabilitation at a single point in time : number of IDUs	Low	Mid	High

## ODT 3 - Outpatient psychosocial support, counselling and psychotherapy

## ODT 3.1: Number of sites providing outpatient psychosocial support, counselling or psychotherapy treatment

Is outpatient psychosocial support, counselling or psychotherapy present?	Yes		
Number of sites - data reported	Psychotherapy, counselling or psychosocial support present but number not known		
Year of reported data	2009		
Reference	World Health Organisation (2009). Global health ATLAS data.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 542 832 null		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Number of outpatient psychosocial support, counselling or psychotherapy sites per 1000 IDUs	Low	Mid	High
	NK	NK	NK

## ODT 3.2: Number of IDUs receiving outpatient psychosocial support, counselling or psychotherapy treatment

Data reported	Psychotherapy, counselling or psychosocial support present but number not known		
Year of reported data	2009		
Reference	World Health Organisation (2009). Global health ATLAS data.		
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)	NK		
Final/Adjusted data for single point in time (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 542 832 null		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Percentage of IDUs in outpatient psychosocial support, counselling or psychotherapy in a 12 month period	Low	Mid	High
	NK	NK	NK
INDICATOR: Percentage of IDUs receiving outpatient psychosocial support, counselling or psychotherapy at a single point in time	Low	Mid	High
	NK	NK	NK

### ODT 3.3: Number of all clients receiving outpatient psychosocial support, counselling or psychotherapy treatment

<b>Data reported</b>	Psychotherapy, counselling or psychosocial support present but number not known		
<b>Year of reported data</b>	2009		
<b>Reference</b>	World Health Organisation (2009). Global health ATLAS data.		
<b>Adjustments</b>			
<b>Final/Adjusted data for 12 month period (to be used as indicator numerator)</b>	NK		
<b>Final/Adjusted data for single point in time (to be used as indicator numerator)</b>	NK		
<b>IDU prevalence estimate (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	0.35	0.45	0.57
<b>Reference</b>	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
<b>General population 15-64 years of age at year of reported data</b>	3 542 832 null		
<b>Estimated number of IDUs at year of reported data (to be used as indicator denominator)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	12 500	16 000	20 000
<b>INDICATOR: Ratio of number of all drug users in outpatient psychosocial support, counselling or psychotherapy in a 12 month period : number of IDUs</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	NK	NK	NK
<b>INDICATOR: Ratio of number of all drug users in outpatient psychosocial support, counselling or psychotherapy at a single point in time : number of IDUs</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	NK	NK	NK

## ODT 4 - Peer-based support groups

## ODT 4.1: Number of peer-based support groups sites

Are Peer-based support groups present?	Yes		
Number of sites - data reported	72 NA meetings per week		
Year of reported data	2008		
Reference	Narcotics Anonymous World Services (2009). Quadrennial report for non-governmental organizations - July 2009.		
Adjustments	Assumed each meeting per week occurs at a separate location		
Final/Adjusted data (to be used as indicator numerator)	72		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 535 039 null		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Number of peer-based support groups sites per 1000 IDUs	Low	Mid	High
	3.60	4.50	5.76

## ODT 4.2: Number of IDUs in peer-based support groups

Data reported	Narcotics Anonymous present but number not known		
Year of reported data	2008		
Reference	Narcotics Anonymous World Services (2009). Quadrennial report for non-governmental organizations - July 2009.		
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)	NK		
Final/Adjusted data for single point in time (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 535 039 null		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Percentage of IDUs in peer-based support groups in a 12 month period	Low	Mid	High
	NK	NK	NK
INDICATOR: Percentage of IDUs in peer-based support groups at a single point in time	Low	Mid	High
	NK	NK	NK

## ODT 4.3: Number of all clients in peer-based support groups

Data reported	Narcotics Anonymous present but number not known		
Year of reported data	2008		
Reference	Narcotics Anonymous World Services (2009). Quadrennial report for non-governmental organizations - July 2009.		
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)	NK		
Final/Adjusted data for single point in time (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 535 039 null		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Ratio of number of all drug users in peer-based support groups in a 12 month period : number of IDUs	Low	Mid	High
	NK	NK	NK
INDICATOR: Ratio of number of all drug users in peer-based support groups at a single point in time : number of IDUs	Low	Mid	High
	NK	null	null

## ODT 5 - Drug treatment (treatment type not specified)

## ODT 5.1: Number of drug treatment sites (treatment type not specified)

Type of drug treatment available	In-patient, out-patient and other treatment - type not specified		
Number of sites - data reported	50 outpatient units 50 in patient units 4 other sites (low-threshold agencies and treatment units in prisons)		
Year of reported data	2007		
Reference	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Treatment demand indicator - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/tdi">http://www.emcdda.europa.eu/stats09/tdi</a> .		
Adjustments	Assume outpatient, inpatient and other units are situated at different locations and counts of sites are therefore mutually exclusive		
Final/Adjusted data (to be used as indicator numerator)	104		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 521 948 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Number of drug treatment sites per 1000 IDUs	Low	Mid	High
	5.20	6.50	8.32

## ODT 5.2: Number of IDUs in drug treatment (treatment type not specified)

<b>Data reported</b>	1198 clients entering treatment at 50 outpatient treatment units 1004 clients entering treatment at 50 inpatient treatment units 19 clients entering treatment at 4 other treatment units 62.9% of clients entering treatment injected primary drug		
<b>Year of reported data</b>	2007		
<b>Reference</b>	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Treatment demand indicator - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/tdi">http://www.emcdda.europa.eu/stats09/tdi</a> .		
<b>Adjustments</b>	Number of IDUs in treatment calculated		
<b>Final/Adjusted data for 12 month period (to be used as indicator numerator)</b>	1397		
<b>Final/Adjusted data for single point in time (to be used as indicator numerator)</b>	NK		
<b>IDU prevalence estimate (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	0.35	0.45	0.57
<b>Reference</b>	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." <i>The Lancet</i> 372(9651): 1733-1745.		
<b>General population 15-64 years of age at year of reported data</b>	3 521 948 (source: UN Population Division, 2009)		
<b>Estimated number of IDUs at year of reported data (to be used as indicator denominator)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	12 500	16 000	20 000
<b>INDICATOR: Percentage of IDUs in drug treatment not specified/drug treatment combined in a 12 month period</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	6.98	8.73	11.18
<b>INDICATOR: Percentage of IDUs in drug treatment not specified/drug treatment combined at a single point in time</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	NK	NK	NK

## ODT 5.3: Number of all clients in drug treatment (treatment type not specified)

<b>Data reported</b>	1198 clients entering treatment at 50 outpatient treatment units 1004 clients entering treatment at 50 inpatient treatment units 19 clients entering treatment at 4 other treatment units		
<b>Year of reported data</b>	2007		
<b>Reference</b>	European Monitoring Centre for Drugs and Drug Addiction. (2009). "Treatment demand indicator - Statistical bulletin." Statistical Bulletin 2009, from <a href="http://www.emcdda.europa.eu/stats09/tdi">http://www.emcdda.europa.eu/stats09/tdi</a> .		
<b>Adjustments</b>			
<b>Final/Adjusted data for 12 month period (to be used as indicator numerator)</b>	2221		
<b>Final/Adjusted data for single point in time (to be used as indicator numerator)</b>	NK		
<b>IDU prevalence estimate (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	0.35	0.45	0.57
<b>Reference</b>	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
<b>General population 15-64 years of age at year of reported data</b>	3 521 948 (source: UN Population Division, 2009)		
<b>Estimated number of IDUs at year of reported data (to be used as indicator denominator)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	12 500	16 000	20 000
<b>INDICATOR: Ratio of number of all drug users in drug treatment not specified/drug treatment combined in a 12 month period : number of IDUs</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	0.11	0.14	0.18
<b>INDICATOR: Ratio of number of all drug users in drug treatment not specified/drug treatment combined at a single point in time : number of IDUs</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	NK	NK	NK

## ODT 6 - Drug treatment in prisons

## ODT 6.1: Number of prison drug treatment sites

Type of drug treatment in prisons?	
Number of sites - data reported	
Year of reported data	
Reference	
Adjustments	
Final/Adjusted data	

## ODT 6.2: Number of IDUs in drug treatment in prisons

Data reported	
Year of reported data	
Reference	
Adjustments	
Final/Adjusted data for 12 month period	
Final/Adjusted data for single point in time	

## ODT 6.3: Number of all clients in drug treatment in prisons

Data reported	
Year of reported data	
Reference	
Adjustments	
Final/Adjusted data for 12 month period	
Final/Adjusted data for single point in time	

## ODT 7 - Compulsory detention as response to drug use

## ODT 7.1: Number of compulsory detention sites for drug use

Number of sites - data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Number of compulsory detention sites for drug use per 1000 IDUs	Low	Mid	High

## ODT 7.2: Number of IDUs in compulsory detention sites for drug use

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)			
Final/Adjusted data for single point in time (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Percentage of IDUs in compulsory detention in closed settings in a 12 month period	Low	Mid	High
INDICATOR: Percentage of IDUs in compulsory detention in closed settings at a single point in time	Low	Mid	High

## ODT 7.3: Number of all drug users in compulsory detention sites for drug use

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data for 12 month period (to be used as indicator numerator)			
Final/Adjusted data for single point in time (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Ratio of all drug users in compulsory detention in closed settings in a 12 month period : number of IDUs	Low	Mid	High
INDICATOR: Ratio of all drug users in compulsory detention in closed settings at a single point in time : number of IDUs	Low	Mid	High

## ODT: All documents reviewed

Bremen Institute for Drug Research (BISDRO) University of Bremen, Centre for Interdisciplinary Addiction Research (CIAR) University of Hamburg, et al. (2008). Final report on prevention, treatment, and harm reduction services in prison, on reintegration services on release from prison and methods to monitor/analyse drug use among prisoners.

European monitoring Centre for Drugs and Drug Addiction. (2008). "Treatment demand indicator - Statistical bulletin " Statistical Bulletin Retrieved 6 April, 2009, from [http://www.emcdda.europa.eu/html.cfm/index65250EN.html?type=stats&stat\\_category=w100&stat\\_type=w87&order=stat\\_reference&CFID=5203606&CFTOKEN=161d43f9a7fd61fb-2D894FE8-E64D-EF3F-72C0BF31D6EAF23A&jsessionid=3830aef6e3d5cdb788212d644b714373741b](http://www.emcdda.europa.eu/html.cfm/index65250EN.html?type=stats&stat_category=w100&stat_type=w87&order=stat_reference&CFID=5203606&CFTOKEN=161d43f9a7fd61fb-2D894FE8-E64D-EF3F-72C0BF31D6EAF23A&jsessionid=3830aef6e3d5cdb788212d644b714373741b).

European Monitoring Centre for Drugs and Drug Addiction. (2009). "Treatment demand indicator - Statistical bulletin." Statistical Bulletin 2009, from <http://www.emcdda.europa.eu/stats09/tdi>.

Helsinki Deaconess Institute (2008). Organization's structure & work

Ronka, S., Virtanen, A., Perala, R. and Vihmo, J. (2007). Finland Drug situation 2007: New developments, trends and in-depth information on selected issues. Helsinki, Finnish National Focal Point, STAKES: 137.

Salminen, M. (2008). UNGASS HIV/AIDS country progress report Finland. Helsinki, KTL - National Public Health Institute, Finland. Department of Infectious Disease Epidemiology and Control. HIV- Unit.

The Helsinki Deaconess Institutes (2008). Information on HIV services for injecting drug users.

World Health Organisation (2009). Global health ATLAS data.

## 4. HIV testing and counselling (T&C)

### T&C 1: Availability of HIV T&C

Is T&C available?	Yes
Most recent year of data	2008
Reference	Salminen, M. (2008). UNGASS HIV/AIDS country progress report Finland. Helsinki, KTL - National Public Health Institute, Finland. Department of Infectious Disease Epidemiology and Control. HIV- Unit.

### T&C 2: Number of sites providing HIV T&C specifically targeting IDUs

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	0	0	0
INDICATOR: Number of targeted T&C sites per 1000 IDUs	Low	Mid	High

### T&C 3: Number of sites providing HIV T&C sites to the general population and also accessible to IDUs

Data reported	VCT available but number not known		
Year of reported data	2008		
Reference	Salminen, M. (2008). UNGASS HIV/AIDS country progress report Finland. Helsinki, KTL - National Public Health Institute, Finland. Department of Infectious Disease Epidemiology and Control. HIV- Unit.		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 535 039 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Number of all T&C sites per 1000 IDUs	Low	Mid	High
	NK	NK	NK

**T&C 4: Number of Individual IDUs who were tested for HIV in 12 month period**

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	0	0	0
INDICATOR: Percentage of individual IDUs accessing T&Cs in a 12 month period	Low	Mid	High

**T&C 5: Percentage of IDUs who were tested for HIV in a 12 month period and who know the result**

Year of reported data	
Reference	
Data reported for both male and female IDUs	
Adjustments	
Final/Adjusted data	
Data reported for male IDUs	
Adjustments	
Final/Adjusted data	
Data reported for female IDUs	
Adjustments	
Final/Adjusted data	

**T&C: All documents reviewed**

Bremen Institute for Drug Research (BISDRO) University of Bremen, Centre for Interdisciplinary Addiction Research (CIAR) University of Hamburg, et al. (2008). Final report on prevention, treatment, and harm reduction services in prison, on reintegration services on release from prison and methods to monitor/analyse drug use among prisoners.

Finnish National Focal Point - STAKES (2008). Finland Drug Situation - New Development, Trends and in-depth information on selected issues. 2008 National report (2007 data) to the EMCDDA. Helsinki.

Helsinki Deaconess Institute (2008). Organization's structure & work

Salminen, M. (2008). UNGASS HIV/AIDS country progress report Finland. Helsinki, KTL - National Public Health Institute, Finland. Department of Infectious Disease Epidemiology and Control. HIV- Unit.

The Helsinki Deaconess Institutes (2008). Information on HIV services for injecting drug users.

**5. Anti-retro-viral (ARV) treatment****ARV 1: Number of sites providing ARV treatment**

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
HIV among IDU prevalence estimate (%)	Low	Mid	High
Reference			
Estimated number of HIV positive IDUs at year of reported data (to be used as indicator)	Low	Mid	High
INDICATOR: Number of ARV treatment sites per 1000 IDUs	Low	Mid	High
INDICATOR: Number of ARV treatment sites per 1000 IDUs living with HIV	Low	Mid	High

## ARV 2: Number of IDUs receiving ARVs at a single point in time

<b>Data reported</b>	100 IDUs on HAART (Transmission route IDU) IDUs represent 11% of people on HAART with known transmission route 35 IDUs on HAART also receiving OST		
<b>Year of reported data</b>	2004		
<b>Reference</b>	Donoghoe, M. C., A. R. Bollerup, et al. (2007). "Access to highly active antiretroviral therapy (HAART) for injecting drug users in the WHO European Region 2002-2004." <i>International Journal of Drug Policy</i> 18(4): 271-80.  WHO and UNAIDS (2008). Progress on implementing the Dublin Declaration on partnership to fight HIV/AIDS in Europe & Central Asia WHO.		
<b>Adjustments</b>			
<b>Final/Adjusted data (to be used as indicator numerator)</b>	100		
<b>IDU prevalence estimate (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	0.35	0.45	0.57
<b>Reference</b>	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." <i>The Lancet</i> 372(9651): 1733-1745.		
<b>General population 15-64 years of age at year of reported data</b>	3 487 624 (source: UN Population Division, 2009)		
<b>Estimated number of IDUs at year of reported data (to be used as indicator denominator)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	12 000	15 500	19 500
<b>HIV among IDU prevalence estimate (%)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	0.10	0.20	0.27
<b>Reference</b>	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." <i>The Lancet</i> 372(9651): 1733-1745.  Upper and lower bounds of range are estimates only, these were calculated as per the methods outlined in Mathers et al 2008.		
<b>Estimated number of HIV positive IDUs at year of reported data (to be used as indicator denominator)</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	12	31	53
<b>INDICATOR: Ratio of IDUs on ART: 100 IDUs living with HIV</b>	<b>Low</b>	<b>Mid</b>	<b>High</b>
	188.93	322.58	830.81

## ARV 3: Restrictions on IDUs' access to ARVs

<b>Do restrictions exist which limit IDUs' access to ARV treatment?</b>	No restrictions
<b>Year of reported data</b>	2008
<b>Reference</b>	Salminen, M. (2008). UNGASS HIV/AIDS country progress report Finland. Helsinki, KTL - National Public Health Institute, Finland. Department of Infectious Disease Epidemiology and Control. HIV- Unit.

**ARV: All documents reviewed**

Bremen Institute for Drug Research (BISDRO) University of Bremen, Centre for Interdisciplinary Addiction Research (CIAR) University of Hamburg, et al. (2008). Final report on prevention, treatment, and harm reduction services in prison, on reintegration services on release from prison and methods to monitor/analyse drug use among prisoners.

Donoghoe, M. C., A. R. Bollerup, et al. (2007). "Access to highly active antiretroviral therapy (HAART) for injecting drug users in the WHO European Region 2002-2004." *International Journal of Drug Policy* 18(4): 271-80.

Salminen, M. (2008). UNGASS HIV/AIDS country progress report Finland. Helsinki, KTL - National Public Health Institute, Finland. Department of Infectious Disease Epidemiology and Control. HIV- Unit.

The Helsinki Deaconess Institutes (2008). Information on HIV services for injecting drug users.

WHO and UNAIDS (2008). Progress on implementing the Dublin Declaration on partnership to fight HIV/AIDS in Europe & Central Asia WHO.

## 6. Condoms

### Condoms 1: Availability of condom programs targeting IDUs

Are condom programs targeting IDUs available?	Yes
Comments	Condoms are distributed at Low Threshold Health Service Centers
Year	2008
Reference	Salminen, M. (2008). UNGASS HIV/AIDS country progress report Finland. Helsinki, KTL - National Public Health Institute, Finland. Department of Infectious Disease Epidemiology and Control. HIV- Unit.

### Condoms 2: Number of sites distributing condoms to IDUs

Data reported	38 health care counselling centres targeting drug users distribute condoms  The Finnish Association of Drug Abusers (Lumme) and the Helsinki Deaconess Institute also distribute condoms but no detail on number of additional sites these represent		
Year of reported data	2008		
Reference	Finnish National Focal Point - STAKES (2008). Finland Drug Situation - New Development, Trends and in-depth information on selected issues. 2008 National report (2007 data) to the EMCDDA. Helsinki.  The Helsinki Deaconess Institutes (2008). Information on HIV services for injecting drug users.  Ronka, S., Virtanen, A., Perala, R. and Vihmo, J. (2007). Finland Drug situation 2007: New developments, trends and in-depth information on selected issues. Helsinki, Finnish National Focal Point, STAKES: 137.		
Adjustments	All NSP sites may distribute condoms but no documentation identified to confirm this		
Final/Adjusted data (to be used as indicator numerator)	38		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 535 039 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Number of condom distribution sites targeting IDU per 1000 IDUs	Low	Mid	High
	1.90	2.38	3.04

## Condoms 3: Number of individual IDUs receiving condoms

Data reported			
Year of reported data			
Reference			
Adjustments			
Final/Adjusted data (to be used as indicator numerator)			
IDU prevalence estimate (%)	Low	Mid	High
Reference			
General population 15-64 years of age at year of reported data			
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
INDICATOR: Percentage of IDUs receiving condoms in a 12 month period	Low	Mid	High

## Condoms 4: Number of condoms distributed to IDUs

Data reported	Condom programs for IDUs present but number not known		
Year of reported data	2008		
Reference	<p>Finnish National Focal Point - STAKES (2008). Finland Drug Situation - New Development, Trends and in-depth information on selected issues. 2008 National report (2007 data) to the EMCDDA. Helsinki.</p> <p>The Helsinki Deaconess Institutes (2008). Information on HIV services for injecting drug users.</p> <p>Ronka, S., Virtanen, A., Perala, R. and Vihmo, J. (2007). Finland Drug situation 2007: New developments, trends and in-depth information on selected issues. Helsinki, Finnish National Focal Point, STAKES: 137.</p>		
Adjustments			
Final/Adjusted data (to be used as indicator numerator)	NK		
IDU prevalence estimate (%)	Low	Mid	High
	0.35	0.45	0.57
Reference	Mathers, B., L. Degenhardt, et al. (2008). "Global epidemiology of injecting drug use and HIV among people who inject drugs: a systematic review." The Lancet 372(9651): 1733-1745.		
General population 15-64 years of age at year of reported data	3 535 039 (source: UN Population Division, 2009)		
Estimated number of IDUs at year of reported data (to be used as indicator denominator)	Low	Mid	High
	12 500	16 000	20 000
INDICATOR: Total number of condoms distributed per 12 month period per IDU	Low	Mid	High
	NK	NK	NK

## Condoms: All documents reviewed

Bremen Institute for Drug Research (BISDRO) University of Bremen, Centre for Interdisciplinary Addiction Research (CIAR) University of Hamburg, et al. (2008). Final report on prevention, treatment, and harm reduction services in prison, on reintegration services on release from prison and methods to monitor/analyse drug use among prisoners.

Finnish National Focal Point - STAKES (2008). Finland Drug Situation - New Development, Trends and in-depth information on selected issues. 2008 National report (2007 data) to the EMCDDA. Helsinki.

Ronka, S., Virtanen, A., Perala, R. and Vihmo, J. (2007). Finland Drug situation 2007: New developments, trends and in-depth information on selected issues. Helsinki, Finnish National Focal Point, STAKES: 137.

Salminen, M. (2008). UNGASS HIV/AIDS country progress report Finland. Helsinki, KTL - National Public Health Institute, Finland. Department of Infectious Disease Epidemiology and Control. HIV- Unit.

The Helsinki Deaconess Institutes (2008). Information on HIV services for injecting drug users.